

JOURNAL

de Théorie des Nombres
de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

János Barát, Peter P. Varjú

A contribution to infinite disjoint covering systems

Tome 17, n° 1 (2005), p. 51-55.

<http://jtnb.cedram.org/item?id=JTNB_2005__17_1_51_0>

© Université Bordeaux 1, 2005, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

A contribution to infinite disjoint covering systems

par JÁNOS BARÁT et PÉTER P. VARJÚ

RÉSUMÉ. Supposons que la famille de suites arithmétiques $\{d_i n + b_i : n \in \mathbb{Z}\}_{i \in I}$ soit un recouvrement disjoint des nombres entiers. Nous prouvons que si $d_i = p^k q^l$ pour des nombres premiers p, q et des entiers $k, l \geq 0$, il existe alors un $j \neq i$ tel que $d_i | d_j$. On conjecture que le résultat de divisibilité est vrai quel que soient les raisons d_i .

Un recouvrement disjoint est appelé saturé si la somme des inverses des raisons est égale à 1. La conjecture ci-dessus est vraie pour des recouvrements saturés avec des d_i dont le produit des facteurs premiers n'est pas supérieur à 1254.

ABSTRACT. Let the collection of arithmetic sequences $\{d_i n + b_i : n \in \mathbb{Z}\}_{i \in I}$ be a disjoint covering system of the integers. We prove that if $d_i = p^k q^l$ for some primes p, q and integers $k, l \geq 0$, then there is a $j \neq i$ such that $d_i | d_j$. We conjecture that the divisibility result holds for all moduli.

A disjoint covering system is called saturated if the sum of the reciprocals of the moduli is equal to 1. The above conjecture holds for saturated systems with d_i such that the product of its prime factors is at most 1254.

A *Beatty sequence* is defined by $S(\alpha, \beta) := \{\lfloor \alpha n + \beta \rfloor\}_{n=1}^{\infty}$, where α is positive and β is an arbitrary real constant. A conjecture of Fraenkel asserts that if $\{S(\alpha_i, \beta_i) : i = 1 \dots m\}$ is a collection of $m \geq 3$ Beatty sequences which partitions the positive integers then α_i / α_j is an integer for some $i \neq j$. Special cases of the conjecture were verified by Fraenkel [2], Graham [4] and Simpson [9]. For more references on Beatty sequences see [1] and [11]. If α is integral, then $S(\alpha, \beta)$ is an arithmetic sequence. Mirsky and Newman, and later independently Davenport and Rado proved that if $\{\{a_i n + b_i\} : i = 1 \dots m\}$ is a partition of the positive integers, then $a_i = a_j$ for some $i \neq j$. This settles Fraenkel's conjecture for integral α 's. We formulate a related conjecture for partitions to infinite number of arithmetic sequences.

We denote by $A(d, b)$ the arithmetic sequence $\{dn + b : n \in \mathbb{Z}\}$. Let a collection \mathcal{S} of arithmetic sequences $\{A(d_i, b_i) : i \in I\}$ be called a *covering*

system (CS), if the union of the sequences is \mathbb{Z} . The CS is finite or infinite according to the set I . The numbers d_i are called the *moduli* of the CS. A conjecture similar to Fraenkel's was posed by Schinzel, that for any finite CS, there is a pair of distinct indices i, j for which $d_i | d_j$. This was verified by Porubský [8] assuming some extra conditions.

When the sequences of a CS are disjoint, it is called a *disjoint covering system* (DCS). The structure of DCS's is a wide topic of research. We only mention here a few results about IIDCS's (such DCS's that the number of sequences are infinite, and the moduli are distinct). For further references see [7]. There is a natural method to construct DCS's, the following construction appeared in [10]:

Example 1. Let $I = \mathbb{N}$, and $d_1 | d_2 | d_3 \dots$ be positive integers. Define the b_i 's recursively to be an integer of minimal absolute value not covered by the sequences $A(d_j, b_j)$, ($j < i$).

Indeed this gives a DCS; if $A(d_i, b_i)$ and $A(d_j, b_j)$ ($j < i$) do intersect then $A(d_i, b_i) \subset A(d_j, b_j)$ as $d_j | d_i$, which contradicts the definition of b_i . Also the definition of b_i guarantees that an integer of absolute value n is covered by one of the first $2n + 1$ sequences.

If the sum of the reciprocals of the moduli equals 1, we call the DCS *saturated*. Apparently every finite DCS is saturated, but this property is rather "rare" for IIDCS's. The IIDCS in the example above is saturated only for $d_i = 2^i$. Stein [10] asked whether this is the unique example. Krukenberg [5] answered this in the negative, then Fraenkel and Simpson [3] characterised all IIDCS, whose moduli are of form $2^k 3^l$. Lewis [6] proved that if a prime greater than 3 divides one of the moduli, then the set of all prime divisors of the moduli is infinite.

We formulate the following conjecture:

Conjecture 2. *If $\{A(d_i, b_i) : i \in I\}$ is a DCS, then for all i there exists an index $j \neq i$ such that $d_i | d_j$.*

This conjecture is valid for the above examples, and also valid for those appearing in [3]. It is also known for finite DCS's, being a consequence of Corollary 2 of [8].

We prove the following special case of Conjecture 2.

Theorem 3. *If $\{A(d_i, b_i) : i \in I\}$ is a DCS, and $d_i = p^k q^l$ for some primes p, q and integers $k, l \geq 0$, then $d_i | d_j$ for some $j \neq i$.*

Proof. Let (a, b) denote the greatest common divisor of the integers a, b , and $[a, b]$ their least common multiple. We will use the fact, that the sequences $A(d_1, b_1)$ and $A(d_2, b_2)$ are disjoint if and only if $(d_1, d_2) \nmid x_1 - x_2$, where x_i is an arbitrary number covered by $A(d_i, b_i)$ for $i = 1, 2$.

If $l = 0$, consider the sequence $A(d_j, b_j)$ that covers $b_i + p^{k-1}$. Then $(d_j, d_i) \nmid b_i + p^{k-1} - b_i = p^{k-1}$. Thus $(d_j, d_i) = p^k = d_i$, so $d_i | d_j$ which was to be proved. Now we may assume that $k, l > 0$.

Assume to the contrary, that the theorem is false. Defining $b'_j = b_j - b_i$, we get another DCS $\{A(d_j, b'_j) : j \in I\}$, where $b'_i = 0$. Hence we may assume that $b_i = 0$.

Let $A_j = A(d_j, b_j) \cap A(p^{k-1}q^{l-1}, 0)$. Either A_j is empty or an arithmetic sequence, whose modulus is $[p^{k-1}q^{l-1}, d_j]$. Let $B_j = \left\{ \frac{x}{p^{k-1}q^{l-1}} \mid x \in A_j \right\}$. The nonempty sequences among the B_j 's form a DCS. Notice that pq divides the modulus of B_j if and only if $d_i = p^k q^l | d_j$. Since the modulus of B_i is pq , it remains to prove the theorem for $k = l = 1$.

Assume $d_i = pq$, and $pq \nmid d_j$ for $i \neq j$. Assume that $p + q$ is covered by the sequence $A(d_t, b_t)$ of the DCS. We prove that $p \nmid d_t$.

Assume to the contrary that $p | d_t$. Let $d_t = d \cdot p^m$, where $p \nmid d$. Then $(pq, d) | q$, and there exist a pair of positive integers u, v such that $q = pq \cdot u - d \cdot v$. Let $a = p + q + dv = p + pqu$. Assume that a is covered by $A(d_s, b_s)$. If $A(d_s, b_s)$ and $A(d_t, b_t)$ are the same sequences, then $d_s = d_t$, and $p | d_s$. Otherwise $(d_s, d_t) \nmid p + q + dv - p - q = dv$, which yields $p | d_s$. Since $pq \nmid a$, $s \neq i$, and $(d_s, d_i) \nmid p + pqu - pqu = p$, thus $q | d_s$. This contradicts $d_i \nmid d_s$.

Similar argument shows $q \nmid d_t$, which contradicts $(d_i, d_t) \nmid b_i - b_t$. So the proof is complete. \square

As a byproduct of the previous proof, we got the following lemma:

Lemma 4. *Suppose there is a DCS $\{A(d_i, b_i) : i \in I\}$ and an index $i \in I$ such, that $d_i = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_k^{\alpha_k}$, where the p 's are distinct primes, and $d_i \nmid d_j$ for all $j \neq i$. Then there exists another DCS $\{A(\hat{d}_i, \hat{b}_i) : i \in \hat{I}\}$ and an index $\hat{i} \in \hat{I}$ such that $\hat{d}_{\hat{i}} = p_1 p_2 \dots p_k$ and $\hat{d}_{\hat{i}} \nmid \hat{d}_j$ for all $j \neq \hat{i}$.*

So it is sufficient to verify the conjecture for square-free moduli. When d_i has more than two different prime factors, the situation seems to be much more complicated. We can still say something for saturated DCS's. We need the following concepts:

Suppose $A \subseteq \mathbb{Z}$. Let $S_n(A) = |\{x \in A : -n < x < n\}|$ be the number of elements of A with absolute value less than n . We define the *density* of A to be $d(A) = \lim_{n \rightarrow \infty} \frac{S_n(A)}{2n-1}$ if the limit exists, and in that case we say, that the density of A exists. We will use following facts. The density is finitely additive, and the density of arithmetic sequences exist, and $d(A(d, b)) = \frac{1}{d}$. Let $\{A(d_i, b_i) : i \in I\}$ be a saturated DCS, and $J \subseteq I$. Lemma 2.2 of [6] states, that the density of $X := \bigcup_{j \in J} A(d_j, b_j)$ exists, and $d(X) = \sum_{j \in J} \frac{1}{d_j}$.

Let a be an arbitrary and b a positive integer. Denote by $a \bmod b$ the unique integer $0 \leq c < b$, that $b|a - c$.

Lemma 5. *Suppose there is a saturated DCS $\{A(d_i, b_i) : i \in I\}$, and an index $i \in I$ such that for all $j \neq i$ $d_i \nmid d_j$. Let D denote the set of positive divisors of d_i different from 1 and d_i . Then there exist nonnegative real numbers $x_{s,t}$, where $s \in D$ and $0 \leq t < s$, such that for all $0 < u < d_i$*

$$\sum_{s \in D} x_{s, u \bmod s} = 1,$$

and $x_{s,0} = 0$ for all $s \in D$.

Proof. Assume $b_i = 0$. Denote by $I_{s,t}$ the set of indices $j \neq i$ such that $(d_i, d_j) = s$, and $b_j \bmod s = t$. Notice that $I_{s,0} = \emptyset$ and $I = \{i\} \cup \bigcup_{s \in D; 0 \leq t < s} I_{s,t}$. Let $y_{s,t} = d(\bigcup_{j \in I_{s,t}} A(d_j, b_j))$. Then $y_{s,0} = 0$ for all s . Since the DCS is saturated, $y_{s,t} = \sum_{j \in I_{s,t}} \frac{1}{d_j}$.

Let $0 < u < d_i$ and

$$Y_{s,u} = \bigcup_{j \in I_{s, u \bmod s}} (A(d_j, b_j) \cap A(d_i, u)).$$

If some element of $A(d_i, u)$ is covered by the sequence $A(d_j, b_j)$, and $(d_i, d_j) = s$, then $s|u - b_j$. Thus $A(d_i, u) = \bigcup_{s \in D} Y_{s,u}$. Notice that $Y_{s,u}$

is the union of sequences of form $A(d_j \cdot \frac{d_i}{s}, u + kd_i)$ for some k , depending on j . Consider the saturated DCS, that consists of the sequences $A(d_j, b_j) \cap A(d_i, u)$, for $j \in I$, $0 \leq u < d_i$. As $Y_{s,u}$ is the union of some sequences of this DCS:

$$d(Y_{s,u}) = \frac{s}{d_i} \sum_{j \in I_{s, u \bmod s}} \frac{1}{d_j} = \frac{s}{d_i} y_{s, u \bmod s}.$$

Since $A(d_i, u) = \bigcup_{s \in D} Y_{s,u}$, $d(d_i, u) = \sum_{s \in D} d(Y_{s,u})$, we get $\frac{1}{d_i} = \sum_{s \in D} \frac{s}{d_i} y_{s, u \bmod s}$.

We finish the proof by setting $x_{s,t} = \frac{y_{s,t}}{s}$. \square

We conjecture, that the system of linear equations in Lemma 5 has no solutions, which would prove Conjecture 2 for saturated DCS's. We have checked this with a computer program for square-free numbers $d_i \leq 1254 = 2 \cdot 3 \cdot 11 \cdot 19$.

Acknowledgements

The authors gratefully thank the referee for her or his useful comments, and Anna Pósfai for the french translation of the abstract. This research has been supported by a Hungarian National Grant, OTKA T.34475.

References

- [1] J. BARÁT, P.P. VARJÚ, *Partitioning the positive integers to seven Beatty sequences*. Indag. Math. **14** (2003), 149–161.
- [2] A.S. FRAENKEL, *Complementing and exactly covering sequences*. J. Combin. Theory Ser. A **14** (1973), 8–20.
- [3] A.S. FRAENKEL, R.J. SIMPSON, *On infinite disjoint covering systems*. Proc. Amer. Math. Soc. **119** (1993), 5–9.
- [4] R.L. GRAHAM, *Covering the positive integers by disjoint sets of the form $\{[n\alpha + \beta] : n = 1, 2, \dots\}$* . J. Combin Theory Ser. A **15** (1973), 354–358.
- [5] C.E. KRUKENBERG, *Covering sets of the integers*. Ph. D. Thesis, Univ. of Illinois, Urbana-Champaign, IL, (1971)
- [6] E. LEWIS, *Infinite covering systems of congruences which don't exist*. Proc. Amer. Math. Soc. **124** (1996), 355–360.
- [7] Š. PORUBSKÝ, J. SCHÖNHEIM, *Covering Systems of Paul Erdős, Past, Present and Future*. In Paul Erdős and his Mathematics I., Springer, Budapest, (2002), 581–627.
- [8] Š. PORUBSKÝ, *Covering systems and generating functions*. Acta Arithm. **26** (1975), 223–231.
- [9] R.J. SIMPSON, *Disjoint covering systems of rational Beatty sequences*. Discrete Math. **92** (1991), 361–369.
- [10] S.K. STEIN, *Unions of Arithmetic Sequences*. Math. Annalen **134** (1958), 289–294.
- [11] R. TIJDEMAN, *Fraenkel's conjecture for six sequences*. Discrete Math. **222** (2000), 223–234.

János BARÁT
Bolyai Institute
University of Szeged
Aradi vértanúk tere 1.
Szeged, 6720 Hungary
E-mail : jbarat@math.u-szeged.hu

Péter VARJÚ
Bolyai Institute
University of Szeged
Aradi vértanúk tere 1.
Szeged, 6720 Hungary
E-mail : Varju.Peter.Pal@stud.u-szeged.hu