

JOURNAL

de Théorie des Nombres
de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Roman URBAN

Sequences of algebraic integers and density modulo 1

Tome 19, n° 3 (2007), p. 755-762.

<http://jtnb.cedram.org/item?id=JTNB_2007__19_3_755_0>

© Université Bordeaux 1, 2007, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

Sequences of algebraic integers and density modulo 1

par ROMAN URBAN

RÉSUMÉ. Nous établissons la densité modulo 1 des ensembles de la forme

$$\{\mu^m \lambda^n \xi + r_m : n, m \in \mathbb{N}\},$$

où $\lambda, \mu \in \mathbb{R}$ sont deux entiers algébriques de degré $d \geq 2$, qui sont rationnellement indépendants et satisfont des hypothèses techniques supplémentaires, $\xi \neq 0$, et r_m une suite quelconque de nombres réels.

ABSTRACT. We prove density modulo 1 of the sets of the form

$$\{\mu^m \lambda^n \xi + r_m : n, m \in \mathbb{N}\},$$

where $\lambda, \mu \in \mathbb{R}$ is a pair of rationally independent algebraic integers of degree $d \geq 2$, satisfying some additional assumptions, $\xi \neq 0$, and r_m is any sequence of real numbers.

1. Introduction

It is a very well known result in the theory of distribution modulo 1 that for every irrational ξ the sequence $\{n\xi : n \in \mathbb{N}\}$ is dense modulo 1 (and even uniformly distributed modulo 1) [11].

In 1967, in his seminal paper [4], Furstenberg proved the following

Theorem 1.1 (Furstenberg, [4, Theorem IV.1]). *If $p, q > 1$ are rationally independent integers (i.e., they are not both integer powers of the same integer) then for every irrational ξ the set*

$$(1.2) \quad \{p^n q^m \xi : n, m \in \mathbb{N}\}$$

is dense modulo 1.

Manuscrit reçu le 17 août 2006.

Research supported in part by the European Commission Marie Curie Host Fellowship for the Transfer of Knowledge “Harmonic Analysis, Nonlinear Analysis and Probability” MTKD-CT-2004-013389 and by the MNiSW research grant N201 012 31/1020.

Mots clefs. Density modulo 1, algebraic integers, topological dynamics, ID-semigroups.

One possible direction of generalizations is to consider p and q in Theorem 1.1 not necessarily integer. This was done by Berend in [3].

According to [10], Furstenberg conjectured that under the assumptions of Theorem 1.1, the set $\{(p^n + q^m)\xi : n, m \in \mathbb{N}\}$ is dense modulo 1. As far as we know, this conjecture is still open. However, there are some results concerning related questions. For example, B. Kra in [9], proved the following

Theorem 1.3 (Kra, [9, Theorem 1.2 and Corollary 2.2]). *For $i = 1, 2$, let $1 < p_i < q_i$ be two rationally independent integers. Assume that $p_1 \neq p_2$ or $q_1 \neq q_2$. Then, for every $\xi_1, \xi_2 \in \mathbb{R}$ with at least one $\xi_i \notin \mathbb{Q}$, the set*

$$\{p_1^n q_1^m \xi_1 + p_2^n q_2^m \xi_2 : n, m \in \mathbb{N}\}$$

is dense modulo 1.

Furthermore, let r_m be any sequence of real numbers and $\xi \notin \mathbb{Q}$. Then, the set

$$(1.4) \quad \{p_1^n q_1^m \xi + r_m : n, m \in \mathbb{N}\}$$

is dense modulo 1.

Inspired by Berend's result [3], we prove some kind of a generalization of the second part of Theorem 1.3 (some kind of an extension of the first part is given in [15]). Namely, we allow algebraic integers, satisfying some additional assumption, to appear in (1.4) instead of integers, and we prove the following

Theorem 1.5. *Let λ, μ be a pair of rationally independent real algebraic integers of degree $d \geq 2$, with absolute values greater than 1. Let $\lambda_2, \dots, \lambda_d$ denote the conjugates of $\lambda = \lambda_1$. Assume that either λ or μ has the property that for every $n \in \mathbb{N}$, its n -th power is of degree d , and that μ may be expressed in the form $g(\lambda)$, where g is a polynomial with integer coefficients, i.e.,*

$$(1.6) \quad \mu = g(\lambda), \text{ for some } g \in \mathbb{Z}[x].$$

Assume further that

$$(1.7) \quad \text{for each } i = 2, \dots, d, \text{ either } |\lambda_i| > 1 \text{ or } |g(\lambda_i)| > 1,$$

and

$$(1.8) \quad \text{for each } i = 2, \dots, d, |\lambda_i| \neq 1.$$

Then for any non-zero ξ , and any sequence of real numbers r_m , the set

$$(1.9) \quad \{\mu^m \lambda^n \xi + r_m : n, m \in \mathbb{N}\}$$

is dense modulo 1.

As an example illustrating Theorem 1.5 we can consider the following expressions

$$(\sqrt{23} + 1)^n(\sqrt{23} + 2)^m + 2^m\beta \text{ or } (3 + \sqrt{3})^n(\sqrt{3})^m 5 + 7^m\beta, \beta \in \mathbb{R}.$$

Remark. We believe that assumption (1.6) is not necessary to conclude density modulo 1 of the sets of the form (1.9).

Another kind of a generalization of Furstenberg's Theorem 1.1, which we are going to use in the proof of our result, is to consider higher-dimensional analogues. A generalization to a commutative semigroup of non-singular $d \times d$ -matrices with integer coefficients acting by endomorphisms on the d -dimensional torus $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$, and to the commutative semigroups of continuous endomorphisms of other compact abelian groups was given by Berend in [1] and [2], respectively (see Sect. 2.3). Recently some results for non-commutative semigroups of endomorphisms of \mathbb{T}^d have been obtained in [5, 6, 13].

The structure of the paper is as follows. In Sect. 2 we recall some notions and facts from ergodic theory and topological dynamics. Following Berend [1, 2], we recall the definition of an ID-semigroup of endomorphisms of the d -dimensional torus \mathbb{T}^d . Then we state Berend's theorem, [1], which gives conditions that guarantee that a given semigroup of endomorphisms of \mathbb{T}^d is an ID-semigroup. This theorem is crucial for the proof of our main result. Finally in Sect. 3, using some ideas from [9, 3] we prove Theorem 1.5.

Acknowledgements. The author wishes to thank the anonymous referee for remarks that improved the overall presentation of the result.

2. Preliminaries

2.1. Algebraic numbers. We say that $P \in \mathbb{Z}[x]$ is *monic* if the leading coefficient of P is one, and *reduced* if its coefficients are relatively prime. A *real algebraic integer* is any real root of a monic polynomial $P \in \mathbb{Z}[x]$, whereas an *algebraic number* is any root (real or complex) of a (not necessarily monic) non-constant polynomial $P \in \mathbb{Z}[x]$. The *minimal polynomial* of an algebraic number θ is the reduced element Q of $\mathbb{Z}[x]$ of the least degree such that $Q(\theta) = 0$. If θ is an algebraic number, the roots of its minimal polynomial are simple. The *degree* of an algebraic number is the degree of its minimal polynomial.

Let θ be an algebraic integer of degree n and let $P \in \mathbb{Z}[x]$ be the minimal polynomial of θ . The $n - 1$ other distinct (real or complex) roots $\theta_2, \dots, \theta_n$ of P are called *conjugates* of θ .

2.2. Topological transitivity, ergodicity and hyperbolic toral endomorphisms. We start with some basic notions, [12, 7]. We consider a *discrete topological dynamical system* (X, f) given by a compact metric

space X and a continuous map $f : X \rightarrow X$. We say that a topological dynamical system (X, f) (or simply that a map f) is *topologically transitive* if for any two nonempty open sets $U, V \subset X$ there exists $n = n(U, V) \in \mathbb{N}$ such that $f^n(U) \cap V \neq \emptyset$. One can show that f is topologically transitive if for every nonempty open set U in X , $\bigcup_{n \geq 0} f^{-n}(U)$ is dense in X (see [8] for other equivalent definitions). If there exists a point $x \in X$ such that its orbit $\{f^n(x) : n \in \mathbb{N}\}$ is dense in X , then we say that x is a *transitive point*. Under some additional assumptions on X , the map f is topologically transitive if and only if there is a transitive point $x \in X$. Namely, we have the following

Proposition 2.1 ([14]). *If X has no isolated point and f has a transitive point then f is topologically transitive. If X is separable, second category and f is topologically transitive then f has a transitive point.*

Consider a probability space (X, \mathcal{F}, μ) and a continuous transformation $f : X \rightarrow X$. We say that the map f is *measure preserving*, and that μ is *f -invariant*, if for every $A \in \mathcal{F}$ we have $\mu(f^{-1}(A)) = \mu(A)$. Recall that f is said to be *ergodic* if every set A such that $f^{-1}(A) = A$ has measure 0 or 1.

Let L be a *hyperbolic matrix*, that is a $d \times d$ -matrix with integer entries, with non-zero determinant, and without eigenvalues of absolute value 1. Then $L\mathbb{Z}^d \subset \mathbb{Z}^d$, so L determines a map of the d -dimensional torus $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$. Such a map is called a *hyperbolic toral endomorphism*. It is known (see e.g. [12]) that the Haar measure m of \mathbb{T}^d is invariant under surjective continuous homomorphisms. In particular, it is L -invariant. We state two propositions about toral endomorphisms. Their proofs can be found in [12].

Proposition 2.2. *Let $L : \mathbb{T}^d \rightarrow \mathbb{T}^d$ be a hyperbolic toral endomorphism. Then L is ergodic.*

The next proposition gives an elementary and useful relation between ergodicity and topological transitivity.

Proposition 2.3. *Let L be a continuous endomorphism of \mathbb{T}^d which preserves the Haar measure m . If L is ergodic then it is topologically transitive. In particular, if L is a hyperbolic toral endomorphism then L has a transitive point $t \in \mathbb{T}^d$, i.e., $\{L^n t : n \in \mathbb{N}\}$ is dense in \mathbb{T}^d .*

We will also need the following lemma about finite invariant sets of ergodic endomorphisms. For the proof see [1, Lemma 5.2].

Lemma 2.4. *Let $L : \mathbb{T}^d \rightarrow \mathbb{T}^d$ be an ergodic endomorphism. A finite L -invariant set is necessarily composed of torsion elements only.*

2.3. ID semigroups of endomorphisms acting on \mathbb{T}^d . Following [1, 2], we say that the semigroup Σ of endomorphisms of a compact group

G has the *ID-property* (or simply that Σ is an *ID-semigroup*) if the only infinite closed Σ -invariant subset of G is G itself. (ID-property stands for *infinite invariant is dense*.) A subset $A \subset G$ is said to be Σ -invariant if $\Sigma A \subset A$.

We say, exactly like in the case of real numbers, that two endomorphisms σ and τ are *rationally dependent* if there are integers m and n , not both of which are 0, such that $\sigma^m = \tau^n$, and *rationally independent* otherwise.

Berend in [1] gave necessary and sufficient conditions in arithmetical terms for a commutative semigroup Σ of endomorphisms of the d -dimensional torus $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$ to have the ID-property. Namely, he proved the following.

Theorem 2.5 (Berend, [1, Theorem 2.1]). *A commutative semigroup Σ of continuous endomorphisms of \mathbb{T}^d has the ID-property if and only if the following hold:*

- (i) *There exists an endomorphism $\sigma \in \Sigma$ such that the characteristic polynomial f_{σ^n} of σ^n is irreducible over \mathbb{Z} for every positive integer n .*
- (ii) *For every common eigenvector v of Σ there exists an endomorphism $\sigma_v \in \Sigma$ whose eigenvalue in the direction of v is of norm greater than 1.*
- (iii) *Σ contains a pair of rationally independent endomorphisms.*

Remark. Let Σ be a commutative ID-semigroup of endomorphisms of \mathbb{T}^d . Then the Σ -orbit of the point $x \in \mathbb{T}^d$ is finite if and only if x is a rational element, i.e., $x = r/q$, $r \in \mathbb{Z}^d$, $q \in \mathbb{N}$ (see [1]).

3. Proof of Theorem 1.5

Let $\lambda > 1$ be a real algebraic integer of degree d with minimal (monic) polynomial $Q_\lambda \in \mathbb{Z}[x]$,

$$Q_\lambda(x) = x^d + c_{d-1}x^{d-1} + \dots + c_1x + c_0.$$

We associate with λ the following *companion matrix* of Q_λ ,

$$\sigma_\lambda = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & & & & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ -c_0 & -c_1 & -c_2 & \dots & -c_{d-1} \end{pmatrix}.$$

Remark. We can think of σ_λ as a matrix of multiplication by λ in the algebraic number field $\mathbb{Q}(\lambda)$. Namely, if $x \in \mathbb{Q}(\lambda)$ has coordinates $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_{d-1})$ in the basis consisting of $1, \lambda, \dots, \lambda^{d-1}$, then λx has coordinates $\alpha \sigma_\lambda$.

Let $\mu = g(\lambda)$, where $g \in \mathbb{Z}[x]$ is a polynomial with integer coefficients, and define the matrix $\sigma_\mu = g(\sigma_\lambda)$.

Denote by Σ the semigroup of endomorphisms of \mathbb{T}^d generated by σ_λ and σ_μ . The vector $v = (1, \lambda, \lambda^2, \dots, \lambda^{d-1})^t$ is an eigenvector of the matrix σ_λ with an eigenvalue λ , that is $\sigma_\lambda v = \lambda v$. Since Σ is a commutative semigroup, it follows that v is a common eigenvector of Σ , in particular $\sigma_\mu v = g(\sigma_\lambda)v = g(\lambda)v = \mu v$.

Clearly, under the assumptions on λ and μ , the operators σ_λ and σ_μ are rationally independent endomorphisms of \mathbb{T}^d and the characteristic polynomial either of σ_λ^n or σ_μ^n is irreducible over \mathbb{Z} for every $n \in \mathbb{N}$. Furthermore, it follows from (1.7) that the condition (ii) of Theorem 2.5 is also satisfied. Thus we have proved the following

Lemma 3.1. *Let λ and μ be as in Theorem 1.5. Let Σ be the semigroup of endomorphisms of \mathbb{T}^d generated by σ_λ and σ_μ . Then Σ is the ID-semigroup.*

The next lemma is a generalization of [9, Lemma 2.1] to the higher-dimensional case. Let X be a compact metric space with a distance d . Consider the space \mathcal{C}_X of all closed subsets of X . The Hausdorff metric d_H on the space \mathcal{C}_X is defined as

$$d_H(A, B) = \max\{\max_{x \in A} d(x, B), \max_{x \in B} d(x, A)\},$$

where $d(x, B) = \min_{y \in B} d(x, y)$ is the distance of x from the set B . It is known that if X is a compact metric space then \mathcal{C}_X is also compact.

Lemma 3.2. *Let σ, τ be a pair of rationally independent and commuting endomorphisms of \mathbb{T}^d . Assume that the semigroup $\Sigma = \langle \sigma, \tau \rangle$ generated by σ and τ satisfies the conditions of Theorem 2.5, and σ is a hyperbolic toral endomorphism of \mathbb{T}^d . Let A be an infinite σ -invariant subset of \mathbb{T}^d . Then for every $\varepsilon > 0$ there exists $m \in \mathbb{N}$ such that the set $\tau^m A$ is ε -dense.*

Proof. It is clear that, taking the closure of A if necessary, we can assume that A is closed. We consider the space $\mathcal{C}_{\mathbb{T}^d}$ of all closed subsets of \mathbb{T}^d with the Hausdorff metric d_H . Let

$$\mathcal{F} := \overline{\{\tau^n A : n \in \mathbb{N}\}} \subset \mathcal{C}_{\mathbb{T}^d}.$$

Since the set A is σ -invariant, it follows that every element (set) $F \in \mathcal{F}$ is also σ -invariant. Define,

$$T = \bigcup_{F \in \mathcal{F}} F \subset \mathbb{T}^d.$$

Since A is an infinite set and $A \subset T$, it follows that T is infinite. Notice that T is closed in \mathbb{T}^d , since \mathcal{F} is closed in $\mathcal{C}_{\mathbb{T}^d}$. Moreover, T is σ - and τ -invariant. Hence, by Theorem 2.5, we get

$$T = \mathbb{T}^d.$$

Since σ is a hyperbolic toral endomorphism, it follows by Proposition 2.3, that there exists $t \in T$ such that the orbit $\{\sigma^{nt} : n \in \mathbb{N}\}$ is dense in \mathbb{T}^d , i.e.,

$$(3.3) \quad \overline{\{\sigma^{nt} : n \in \mathbb{N}\}} = \mathbb{T}^d$$

Clearly, $t \in F$ for some $F \in \mathcal{F}$. By definition of \mathcal{F} , there is a sequence $\{n_k\} \subset \mathbb{N}$ such that $F = \lim_k \tau^{n_k} A$, and the limit is taken in the Hausdorff metric d_H . Since $t \in F$ and F is σ -invariant, we get $F \supset \overline{\{\sigma^{nt} : n \in \mathbb{N}\}} = \mathbb{T}^d$ (see (3.3)). Hence, $F = \mathbb{T}^d$. Therefore, for sufficiently large k , $\tau^{n_k} A$ is ε -dense. \square

Now we are ready to give

Proof of Theorem 1.5. Let $\alpha = \xi(1, \lambda, \lambda^2, \dots, \lambda^{d-1})^t \in \mathbb{R}^d$ be a common eigenvector of the semigroup Σ . Consider

$$A = \{\sigma_\lambda^n \pi(\alpha) : n \in \mathbb{N}\} = \{\pi(\lambda^n \xi, \lambda^{n+1} \xi, \dots, \lambda^{n+d-1} \xi)^t : n \in \mathbb{N}\},$$

where $\pi : \mathbb{R}^d \rightarrow \mathbb{T}^d$ is the canonical projection. By (1.8), σ_λ is a hyperbolic toral endomorphism. In particular, by Proposition 2.2, σ_λ is ergodic. Since $\pi(\alpha)$ is not a torsion element, it follows from Lemma 2.4 that A is infinite. By Lemma 3.1, $\Sigma = \langle \sigma_\lambda, \sigma_\mu \rangle$ is the ID-semigroup of \mathbb{T}^d . Thus, by Lemma 3.2 applied to σ_λ and σ_μ , there exists $m \in \mathbb{N}$ such that $\sigma_\mu^m A$ is ε -dense. Let $v_m = \pi(r_m, 0, \dots, 0)^t$. Since

$$\sigma_\mu^m A + v_m = \{\pi(\mu^m \lambda^n \xi + r_m, \mu^m \lambda^{n+1} \xi, \dots, \mu^m \lambda^{n+d-1} \xi)^t : n \in \mathbb{N}\}$$

is a translate of an ε -dense set, it is also ε -dense. Now, taking the projection of the set $\sigma_\mu^m A + v_m$ on the first coordinate we get the result. \square

References

- [1] D. BEREND, *Multi-invariant sets on tori*. Trans. Amer. Math. Soc. **280** (1983), no. 2, 509–532.
- [2] D. BEREND, *Multi-invariant sets on compact abelian groups*. Trans. Amer. Math. Soc. **286** (1984), no. 2, 505–535.
- [3] D. BEREND, *Dense (mod 1) dilated semigroups of algebraic numbers*. J. Number Theory **26** (1987), no. 3, 246–256.
- [4] H. FURSTENBERG, *Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation*. Math. Systems Theory **1** (1967), 1–49.
- [5] Y. GUIVARCH AND A. N. STARKOV, *Orbits of linear group actions, random walk on homogeneous spaces, and toral automorphisms*. Ergodic Theory Dynam. Systems **24** (2004), no. 3, 767–802.
- [6] Y. GUIVARCH AND R. URBAN, *Semigroup actions on tori and stationary measures on projective spaces*. Studia Math. **171** (2005), no. 1, 33–66.
- [7] A. KATOK AND B. HASSELBLATT, *Introduction to the modern theory of dynamical systems*. Encyclopedia of Mathematics and its Applications **54**, Cambridge University Press, Cambridge, 1995.
- [8] S. KOLYADA AND L. SNOHA, *Some aspects of topological transitivity – a survey*. Grazer Math. Ber. **334** (1997), 3–35.
- [9] B. KRA, *A generalization of Furstenberg’s Diophantine theorem*. Proc. Amer. Math. Soc. **127** (1999), no. 7, 1951–1956.

- [10] D. MEIRI, *Entropy and uniform distribution of orbits in \mathbb{T}^d* . Israel J. Math. **105** (1998), 155–183.
- [11] L. KUIPERS AND H. NIEDERREITER, *Uniform distribution of sequences*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1974.
- [12] R. MAÑÉ, *Ergodic theory and differentiable dynamics*. Ergebnisse der Mathematik und ihrer Grenzgebiete **3**. Springer-Verlag, Berlin, 1987.
- [13] R. MUCHNIK, *Semigroup actions on \mathbb{T}^n* . Geometriae Dedicata **110** (2005), 1–47.
- [14] S. SILVERMAN, *On maps with dense orbits and the definition of chaos*. Rocky Mt. J. Math. **22** (1992), no. 1, 353–375.
- [15] R. URBAN, *On density modulo 1 of some expressions containing algebraic integers*. Acta Arith., **127** (2007), no. 3, 217–229.

Roman URBAN
Institute of Mathematics
Wroclaw University
Plac Grunwaldzki 2/4
50-384 Wroclaw, Poland
E-mail: urban@math.uni.wroc.pl