

JOURNAL

de Théorie des Nombres

de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Kevin J. MCGOWN

Norm-Euclidean Galois fields and the Generalized Riemann Hypothesis

Tome 24, n° 2 (2012), p. 425-445.

<http://jtnb.cedram.org/item?id=JTNB_2012__24_2_425_0>

© Société Arithmétique de Bordeaux, 2012, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

Norm-Euclidean Galois fields and the Generalized Riemann Hypothesis

par KEVIN J. MCGOWN

RÉSUMÉ. En supposant que l'hypothèse de Riemann généralisée (HRG) soit vérifiée, nous montrons que les corps de nombres galoisiens de degré 3 qui sont euclidiens pour la norme sont précisément ceux dont le discriminant est l'un des entiers suivants :

$$\Delta = 7^2, 9^2, 13^2, 19^2, 31^2, 37^2, 43^2, 61^2, 67^2, 103^2, 109^2, 127^2, 157^2.$$

Une grande partie de la preuve consiste à établir le résultat plus général suivant : soit K un corps de nombres galoisien de degré premier impair ℓ et de conducteur f . Supposons que HRG soit vérifiée pour $\zeta_K(s)$. Si

$$38(\ell - 1)^2(\log f)^6 \log \log f < f,$$

alors K n'est pas euclidien pour la norme.

ABSTRACT. Assuming the Generalized Riemann Hypothesis (GRH), we show that the norm-Euclidean Galois cubic fields are exactly those with discriminant

$$\Delta = 7^2, 9^2, 13^2, 19^2, 31^2, 37^2, 43^2, 61^2, 67^2, 103^2, 109^2, 127^2, 157^2.$$

A large part of the proof is in establishing the following more general result: Let K be a Galois number field of odd prime degree ℓ and conductor f . Assume the GRH for $\zeta_K(s)$. If

$$38(\ell - 1)^2(\log f)^6 \log \log f < f,$$

then K is not norm-Euclidean.

1. Introduction

Let K be a number field with ring of integers \mathcal{O}_K , and denote by $N = N_{K/\mathbb{Q}}$ the absolute norm map. For brevity, we will sometimes use the term field to mean a number field. We call a number field K norm-Euclidean if for every $\alpha, \beta \in \mathcal{O}_K$, $\beta \neq 0$, there exists $\gamma \in \mathcal{O}_K$ such that $|N(\alpha - \gamma\beta)| < |N(\beta)|$. In the quadratic setting, it is known that there

Manuscrit reçu le 14 février 2011.

Mots clefs. norm-Euclidean, Galois fields, cubic fields, GRH, Dirichlet characters.

Classification math. 11A05, 11R04, 11R16, 11R80, 11Y40.

are only finitely many norm-Euclidean fields and they have been identified; namely, a number field of the form $K = \mathbb{Q}(\sqrt{d})$ with d squarefree is norm-Euclidean if and only if

$$d = -1, -2, -3, -7, -11, 2, 3, 5, 6, 7, 11, 13, 17, 19, 21, 29, 33, 37, 41, 57, 73.$$

The main goal of this paper is to prove the following:

Theorem 1.1. *Assuming the GRH, the norm-Euclidean Galois cubic fields are exactly those with discriminant*

$$\Delta = 7^2, 9^2, 13^2, 19^2, 31^2, 37^2, 43^2, 61^2, 67^2, 103^2, 109^2, 127^2, 157^2.$$

For most of this paper, the reader may take the Generalized Riemann Hypothesis (GRH) to mean that for every Dirichlet L -function $L(s, \chi)$, all the zeros of $L(s, \chi)$ in the critical strip $0 < \Re(s) < 1$ are on the critical line $\Re(s) = 1/2$.¹ The only exceptions will be when we explicitly state which function is being referred to — i.e., “the GRH for $L(s, \chi)$ ” or “the GRH for $\zeta_K(s)$ ”.

Previously, Heilbronn (see [5]) showed that there are finitely many norm-Euclidean Galois cubic fields, but produced no upper bound on the discriminant. Godwin and Smith (see [4]) showed that the Galois cubic fields with $|\Delta| < 10^8$ are exactly those listed in Theorem 1.1 and were the first to give this list. Lemmermeyer subsequently extended this result to show that Godwin and Smith’s list includes all fields with $|\Delta| < 2.5 \cdot 10^{11}$ (see [9]). Although it is the natural question, no one seems to have conjectured that this is the complete list; however, in light of Theorem 1.1, this now seems like a very reasonable conjecture!

In a recent paper (see [10]) the author proved the following unconditional result:

Theorem 1.2. *The fields listed in Theorem 1.1 are norm-Euclidean, and any remaining norm-Euclidean Galois cubic field must have discriminant $\Delta = f^2$ with $f \equiv 1 \pmod{3}$ where f is a prime in the interval $(10^{10}, 10^{70})$.*

A large part of the proof of Theorems 1.1 and 1.2 is in giving an upper bound on the discriminant for the class of fields in question. Our technique works not only in the case of Galois cubic fields, but for Galois fields of odd prime degree.

Theorem 1.3. *Let ℓ be an odd prime. There exists a computable constant C_ℓ such that if K is a Galois number field of odd prime degree ℓ , conductor f , and discriminant Δ , which is norm-Euclidean, then $f < C_\ell$ and $0 < \Delta < C_\ell^{\ell-1}$.*

¹Actually, for Theorem 1.1, it suffices to assume the Riemann Hypothesis (RH) and the GRH for Dirichlet L -functions associated to cubic characters.

ℓ	C_ℓ	ℓ	C_ℓ	ℓ	C_ℓ
3	10^{70}	29	10^{98}	61	10^{106}
5	10^{78}	31	10^{99}	67	10^{107}
7	10^{82}	37	10^{101}	71	10^{107}
11	10^{88}	41	10^{102}	73	10^{108}
13	10^{89}	43	10^{102}	79	10^{108}
17	10^{92}	47	10^{103}	83	10^{109}
19	10^{94}	53	10^{104}	89	10^{109}
23	10^{96}	59	10^{105}	97	10^{110}

TABLE 1.1. Values of C_ℓ for primes $\ell < 100$

ℓ	C_ℓ	ℓ	C_ℓ	ℓ	C_ℓ
3	10^{11}	29	10^{15}	61	10^{15}
5	10^{12}	31	10^{15}	67	10^{15}
7	10^{13}	37	10^{15}	71	10^{16}
11	10^{13}	41	10^{15}	73	10^{16}
13	10^{14}	43	10^{15}	79	10^{16}
17	10^{14}	47	10^{15}	83	10^{16}
19	10^{14}	53	10^{15}	89	10^{16}
23	10^{14}	59	10^{15}	97	10^{16}

TABLE 1.2. Values of C_ℓ for primes $\ell < 100$, assuming the GRH

In [10], the author proved Theorem 1.3 and gave the constants in Table 1.1. In this paper, we show that under the GRH these constants can be improved to those given in Table 1.2. In fact, we prove the following result which, after some easy computation, completely justifies Table 1.2.

Theorem 1.4. *Let K be a Galois number field of odd prime degree ℓ and conductor f . Assume the GRH for $\zeta_K(s)$, the Dedekind zeta function of K .² If*

$$38(\ell - 1)^2(\log f)^6 \log \log f < f,$$

then K is not norm-Euclidean.

2. Preliminaries

As is customary, we write $\zeta(s)$ to denote the Riemann zeta function, $L(s, \chi)$ to denote the Dirichlet L -function associated to a Dirichlet character χ , and $\zeta_K(s)$ to denote the Dedekind zeta function associated to a number field K . The following is well-known and is an easy consequence of Theorem 8.6 of [12].

²Note that, in this context, the GRH (as defined above) implies the GRH for $\zeta_K(s)$; this follows from Lemma 2.1.

Lemma 2.1. *Let K be a Galois number field of odd prime degree ℓ and conductor f , and let χ be a primitive Dirichlet character modulo f of order ℓ . Then*

$$\zeta_K(s) = \zeta(s) \prod_{k=1}^{\ell-1} L(s, \chi^k).$$

We now quote three results from [10] which will be crucial for our arguments.

Theorem 2.1. *Let K be a Galois number field of odd prime degree ℓ , conductor f , and discriminant Δ . If K is norm-Euclidean, then f is a prime with $f \equiv 1 \pmod{\ell}$ and $\Delta = f^{\ell-1}$, except when K is the cubic field with $f = 9$ and $\Delta = 81$.*

Theorem 2.2. *Let K be a Galois number field of odd prime degree ℓ and conductor f with $(f, \ell) = 1$, and let χ be a primitive Dirichlet character modulo f of order ℓ . Denote by $q_1 < q_2$ the two smallest rational primes that are inert in K . Suppose that there exists $r \in \mathbb{Z}^+$ with*

$$(r, q_1 q_2) = 1, \quad \chi(r) = \chi(q_2)^{-1},$$

such that any of the following conditions hold:

- (1) $r q_2 k \not\equiv f \pmod{q_1^2}, \quad k = 1, \dots, q_1 - 1,$
 $(q_1 - 1)(q_2 r - 1) \leq f$
- (2) $q_1 \neq 2, 3, \quad 3q_1 q_2 r \log q_1 < f$
- (3) $q_1 \neq 2, 3, 7, \quad 2.1 q_1 q_2 r \log q_1 < f$
- (4) $q_1 = 2, q_2 \neq 3, \quad 3q_2 r < f$
- (5) $q_1 = 3, q_2 \neq 5, \quad 5q_2 r < f$

Then K is not norm-Euclidean.

Proposition 2.1. *Let K be a Galois number field of odd prime degree ℓ and conductor f . Denote by $q_1 < q_2$ the two smallest rational primes that are inert in K . Suppose either of the following conditions hold:*

- (1) $q_1 = 2, q_2 = 3,$
 $72(\ell - 1)f^{1/2} \log 4f + 35 \leq f$
- (2) $q_1 = 3, q_2 = 5,$
 $507(\ell - 1)f^{1/2} \log 9f + 448 \leq f$

Then K is not norm-Euclidean.

3. GRH bounds for non-residues

In [2], Bach proves an explicit version of a theorem due to Ankeny (see [1]) regarding the least element outside of a given non-trivial subgroup of $(\mathbb{Z}/m\mathbb{Z})^*$. The main idea behind Bach’s proof appears in [11], but to obtain explicit results there are many details to work out; Bach uses a slightly different kernel and introduces a parameter in order to achieve good

numerical results. Using the tables in [2], we obtain the following special case which is useful to us in the present context.

Theorem 3.1 (Bach, 1990). *Assume the GRH. Let χ be a non-principal³ Dirichlet character modulo $m \geq 10^8$, and denote by q_1 the smallest prime such that $\chi(q_1) \neq 1$. Then*

$$q_1 < (1.17 \log m - 6.36)^2.$$

We will follow Bach’s approach to give bounds on the quantities q_2 and r appearing in Theorem 2.2. Although the following results undoubtedly hold in more generality, we will not hesitate to specialize to our situation when it affords us certain technical conveniences.

Theorem 3.2. *Let χ be a non-principal Dirichlet character modulo $m \geq 10^9$ with $\chi(-1) = 1$. Assume the RH and the GRH for $L(s, \chi)$. Denote by $q_1 < q_2$ the two smallest primes such that $\chi(q_1), \chi(q_2) \neq 1$. Then*

$$q_2 < 2.5(\log m)^2.$$

Theorem 3.3. *Let ℓ and f be odd primes with $f \geq 10^8$ and $f \equiv 1 \pmod{\ell}$. Let K be the Galois number field of degree ℓ and conductor f , and let χ be a primitive Dirichlet character modulo f of order ℓ . Assume the GRH for $\zeta_K(s)$. If q_1, q_2 are rational primes and $\omega \neq 1$ is an ℓ -th root of unity, then there exists $r \in \mathbb{Z}^+$ such that $(r, q_1 q_2) = 1$, $\chi(r) = \omega$, and*

$$r < 2.5(\ell - 1)^2(\log f)^2.$$

The remainder of §3 is devoted to proving Theorems 3.2 and 3.3. Although this constitutes the bulk of the paper and is where the analytic techniques come into play, the casual reader who is willing to accept these two results may skip the rest of this section and proceed to §4.

In §3.1 we give some explicit formulas relating sums over prime powers to sums over zeros of L -functions, and in §3.2 we give some GRH estimates for the sums over zeros. Then in §3.3 and §3.4 we prove Theorems 3.2 and 3.3, respectively.

3.1. An explicit formula.

Lemma 3.1. *Let χ be a Dirichlet character modulo m . (Here we allow the possibility that χ is the principal character or even that $m = 1$.) For $x > 1$ and $a \in (0, 1)$, we have*

$$-\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{x^s}{(s+a)^2} \frac{L'(s, \chi)}{L(s, \chi)} ds = \sum_{n < x} \chi(n) \Lambda(n) (n/x)^a \log(x/n).$$

³The principal character modulo m is the Dirichlet character $\chi : \mathbb{Z} \rightarrow \mathbb{C}$ induced by the trivial character on $(\mathbb{Z}/m\mathbb{Z})^*$.

Proof. This is Lemma 4.2 of [2]. We provide only a brief sketch here. We plug the Dirichlet series

$$\frac{L'(s, \chi)}{L(s, \chi)} = - \sum_{n=1}^{\infty} \chi(n)\Lambda(n)n^{-s}$$

into the left-hand side above and interchange the order of summation and integration. Next, we use the fact that for $y > 0$ one has

$$\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{y^s}{(s+a)^2} ds = \begin{cases} y^{-a} \log y & \text{if } y > 1 \\ 0 & \text{otherwise} \end{cases},$$

and the result follows. □

Lemma 3.2. *Let χ be a non-principal primitive Dirichlet character modulo m with $\chi(-1) = 1$. For $x > 1$ and $a \in (0, 1)$ we have*

$$\begin{aligned} \sum_{n < x} \chi(n)\Lambda(n)(n/x)^a \log(x/n) &= - \sum_{\rho \text{ of } L_\chi} \frac{x^\rho}{(\rho+a)^2} - \sum_{n=1}^{\infty} \frac{x^{-2n}}{(a-2n)^2} - \frac{1}{a^2} \\ &\quad - \frac{\log x}{x^a} \left(\frac{L'_\chi}{L_\chi} \right) (-a) - \frac{1}{x^a} \left(\frac{L'_\chi}{L_\chi} \right)' (-a). \end{aligned}$$

Proof. Formally, this follows immediately by evaluating the integral in Lemma 3.1 by residues. For more details, see Lemma 4.4 of [2]. □

Lemma 3.3. *For $x > 1$ and $a \in (0, 1)$ we have*

$$\begin{aligned} \sum_{n < x} \Lambda(n)(n/x)^a \log(x/n) &= \frac{x}{(a+1)^2} - \sum_{\rho \text{ of } \zeta} \frac{x^\rho}{(\rho+a)^2} - \sum_{n=1}^{\infty} \frac{x^{-2n}}{(a-2n)^2} \\ &\quad - \frac{\log x}{x^a} \left(\frac{\zeta'}{\zeta} \right) (-a) - \frac{1}{x^a} \left(\frac{\zeta'}{\zeta} \right)' (-a). \end{aligned}$$

Proof. The proof is similar to the proof of the previous result. □

For our bounds on q_2 and r , we will need to exclude certain primes from consideration; this will require the following estimate:

Lemma 3.4. *Let $u \in \mathbb{Z}^+$. Then*

$$\sum_{\substack{n < x \\ (n,u) > 1}} \Lambda(n)(n/x)^a \log(x/n) \leq \omega(u)(\log x)^2,$$

where $\omega(u)$ denotes the number of distinct prime factors of u .

Proof. If $u = 1$, the result is trivial. Suppose $u = p_1^{a_1} \dots p_t^{a_t}$. Then

$$\sum_{\substack{n < x \\ (n,u) > 1}} \Lambda(n) = \sum_{k=1}^t \sum_{a=1}^{\lfloor \log_{p_k} x \rfloor} \log p_k \leq \sum_{k=1}^t \log x = t \log x.$$

The result easily follows. □

3.2. Sums over zeros. In order to prove our results, we will need to bound the sums over zeros appearing in Lemmas 3.2 and 3.3. Eventually we will take character combinations of the formulas appearing in these lemmas as well, and so it will be useful to bound the corresponding sum over all the zeros of the Dedekind zeta function of a number field K .

Let K be a number field of discriminant Δ with r_1 real embeddings and $2r_2$ complex embeddings. We define

$$\psi_K(s) := \frac{r_1 + r_2}{2} \psi\left(\frac{s}{2}\right) + \frac{r_2}{2} \psi\left(\frac{s+1}{2}\right) - \frac{1}{2} [K : \mathbb{Q}] \log \pi, \quad \psi(s) := \frac{\Gamma'(s)}{\Gamma(s)},$$

where $\Gamma(s)$ is the usual gamma function. In particular,

$$\psi_{\mathbb{Q}}(s) = \frac{1}{2} \left(\psi\left(\frac{s}{2}\right) - \log \pi \right).$$

In order to expedite the proofs of this section, we quote some formulae, all of which can be derived from (5.9) of [7]. For all $s \in \mathbb{C}$, we have:

$$(3.1) \quad \frac{\zeta'_K(s)}{\zeta_K(s)} = B_K + \sum_{\rho \text{ of } \zeta_K} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right) - \frac{1}{2} \log |\Delta| - \frac{1}{s} - \frac{1}{s-1} - \psi_K(s)$$

$$(3.2) \quad \frac{\zeta'(s)}{\zeta(s)} = B + \sum_{\rho \text{ of } \zeta} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right) - \frac{1}{s} - \frac{1}{s-1} - \psi_{\mathbb{Q}}(s)$$

If χ is a non-principal primitive Dirichlet character modulo f , with $\chi(-1) = 1$, then for all $s \in \mathbb{C}$ we have:

$$(3.3) \quad \frac{L'(s, \chi)}{L(s, \chi)} = B_{\chi} + \sum_{\rho \text{ of } L_{\chi}} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right) - \frac{1}{2} \log f - \psi_{\mathbb{Q}}(s)$$

Each sum above is over the non-trivial zeros ρ of the corresponding functions, and is absolutely and uniformly convergent on compact subsets of \mathbb{C} . Henceforth we adopt the notation that ρ will always denote a non-trivial zero with $0 < \Re(\rho) < 1$.

Each of (3.1), (3.2), (3.3) involves a constant B which can be difficult to estimate. Fortunately, in all three cases this constant can be eliminated from the equation as follows. Provided the sum is taken in symmetric order⁴, one has

$$(3.4) \quad B + \sum_{\rho \text{ of } \zeta} \frac{1}{\rho} = 0,$$

⁴Taking the sum in symmetric order means: $\sum_{\rho} = \lim_{T \rightarrow \infty} \sum_{\substack{\rho = \sigma + it \\ |t| < T}}$

and similarly for B_K and B_χ . See [3] for a simple argument which gives this result for the constant B . The corresponding result for B_K follows by a similar argument and was first exploited by Stark to give lower bounds for discriminants (see [16, 17]).

The analogous result for B_χ is not obvious; in fact, it wasn't known until the introduction of the Weil formulas (see [18, 19]). Plugging $s = 1$ into (3.3) and comparing against (2.3.1) of [6] gives a proof of this result. See [14, 15, 13] for results regarding the use of explicit formulae to obtain discriminant bounds.

We begin with a lemma which goes back to Landau (see [8]).

Lemma 3.5. *Let χ be a primitive Dirichlet character modulo f with $\chi(-1) = 1$. Then for $\sigma \in \mathbb{R}$, we have*

$$\sum_{\rho \text{ of } L_\chi} \left(\frac{1}{\sigma - \rho} + \frac{1}{\sigma - \bar{\rho}} \right) = \log f + 2\Re \frac{L'(\sigma, \chi)}{L(\sigma, \chi)} + 2\psi_{\mathbb{Q}}(\sigma).$$

Proof. We substitute $s = \sigma$ into (3.3) and add the result to its conjugate. The result now follows upon invoking the fact that

$$\Re \left(B_\chi + \sum_{\rho \text{ of } L_\chi} \frac{1}{\rho} \right) = 0.$$

□

Lemma 3.6. *Let χ be a non-principal primitive Dirichlet character modulo f with $\chi(-1) = 1$. Assume the RH and the GRH for $L(s, \chi)$. For $a \in (0, 1)$ we have*

$$\sum_{\rho \text{ of } \zeta, L_\chi} \frac{1}{|\rho + a|^2} \leq \frac{1}{2a + 1} \left(\log f + 2 \left(\frac{1}{a + 1} + \frac{1}{a} \right) + 4\psi_{\mathbb{Q}}(a + 1) \right).$$

Proof. We consider the following two formulae:

$$\begin{aligned} \sum_{\rho \text{ of } \zeta} \left(\frac{1}{\sigma - \rho} + \frac{1}{\sigma - \bar{\rho}} \right) &= 2 \frac{\zeta'(\sigma)}{\zeta(\sigma)} + 2 \left(\frac{1}{\sigma} + \frac{1}{\sigma - 1} \right) + 2\psi_{\mathbb{Q}}(\sigma) \\ \sum_{\rho \text{ of } L_\chi} \left(\frac{1}{\sigma - \rho} + \frac{1}{\sigma - \bar{\rho}} \right) &= \log f + 2\Re \frac{L'(\sigma, \chi)}{L(\sigma, \chi)} + 2\psi_{\mathbb{Q}}(\sigma) \end{aligned}$$

The second formula above is Lemma 3.5 and the first can be proved in exactly the same manner. Setting $\sigma = a + 1$ and supposing that $\Re(\rho) = 1/2$, we find:

$$(3.5) \quad \frac{1}{|\rho + a|^2} = \frac{1}{2a + 1} \left(\frac{1}{\sigma - \rho} + \frac{1}{\sigma - \bar{\rho}} \right)$$

To complete the proof, we combine everything above and note that

$$\frac{\zeta'(\sigma)}{\zeta(\sigma)} + \Re \frac{L'(\sigma, \chi)}{L(\sigma, \chi)} < 0,$$

by considering the Dirichlet series for $(\zeta'/\zeta + L'_\chi/L_\chi)(s)$. □

We give a special case of the previous lemma:

Lemma 3.7. *Let χ be a non-principal primitive Dirichlet character modulo f with $\chi(-1) = 1$. Assume the RH and the GRH for $L(s, \chi)$. We have*

$$\sum_{\rho \text{ of } \zeta, L_\chi} \frac{1}{\left| \rho + \frac{1}{2} \right|^2} \leq \frac{1}{2} \log f + 0.437.$$

Proof. Use the fact

$$(3.6) \quad \psi_{\mathbb{Q}}(3/2) \approx -1.1153$$

and apply the previous lemma with $a = 1/2$. □

Having completed the desired estimates over the zeros of $\zeta(s)$ and $L(s, \chi)$, we turn to $\zeta_K(s)$.

Lemma 3.8. *Let K be a number field with discriminant Δ . Then we have*

$$\sum_{\rho \text{ of } \zeta_K} \left(\frac{1}{\sigma - \rho} + \frac{1}{\sigma - \bar{\rho}} \right) = \log |\Delta| + 2 \left(\frac{1}{\sigma} + \frac{1}{\sigma - 1} \right) + 2\psi_K(\sigma) + 2 \frac{\zeta'_K(\sigma)}{\zeta_K(\sigma)}.$$

Proof. This is exactly analogous to Lemma 3.5. □

Lemma 3.9. *Let K be a number field with discriminant Δ . Suppose the GRH holds for $\zeta_K(s)$. For $a \in (0, 1)$ we have*

$$\sum_{\rho \text{ of } \zeta_K} \frac{1}{|\rho + a|^2} < \frac{1}{2a + 1} \left[\log |\Delta| + 2 \left(\frac{1}{a + 1} + \frac{1}{a} \right) + 2\psi_K(a + 1) \right].$$

Proof. Let $\sigma = a + 1$. Apply Lemma 3.8, use (3.5), and observe that $\zeta'_K(\sigma)/\zeta_K(\sigma) < 0$. □

We give a special case of the previous lemma:

Lemma 3.10. *Let K be a totally real number field with discriminant Δ . Suppose the GRH holds for $\zeta_K(s)$. We have*

$$\sum_{\rho \text{ of } \zeta_K} \frac{1}{\left| \rho + \frac{1}{2} \right|^2} < \frac{1}{2} \left(\log |\Delta| + \frac{16}{3} + 2 \psi_{\mathbb{Q}}(3/2) [K : \mathbb{Q}] \right).$$

Proof. Since $r_1 = [K : \mathbb{Q}]$, $r_2 = 0$, we have $\psi_K(s) = [K : \mathbb{Q}] \psi_{\mathbb{Q}}(s)$. The result now follows from the previous lemma upon setting $a = 1/2$. □

Now we specialize even further to our situation:

Lemma 3.11. *Let K be a totally real number field of degree ℓ and discriminant $\Delta = f^{\ell-1}$. Suppose the GRH holds for $\zeta_K(s)$. We have*

$$\sum_{\rho \text{ of } \zeta_K} \frac{1}{\left| \rho + \frac{1}{2} \right|^2} < \frac{1}{2} [(\ell - 1) \log f - 2.23 \ell + 5.34].$$

Proof. We apply the previous lemma, using the approximation given in (3.6). □

3.3. An upper estimate on q_2 . We establish a series of results, building up to the proof of Theorem 3.2.

Lemma 3.12. *Let χ be a non-principal Dirichlet character modulo m with $\chi(-1) = 1$. For $a \in (0, 1)$ and $x > 0$ we have*

$$\begin{aligned} \frac{x}{(a+1)^2} + \frac{1}{a^2} &= \sum_{\rho \text{ of } \zeta} \frac{x^\rho}{(\rho+a)^2} - \sum_{\rho \text{ of } L_\chi} \frac{x^\rho}{(\rho+a)^2} \\ &+ \sum_{\substack{n < x \\ \chi(n) \neq 1}} (1 - \chi(n)) \Lambda(n) (n/x)^a \log(x/n) \\ &+ \frac{\log x}{x^a} \left[\left(\frac{\zeta'}{\zeta} \right) (-a) - \left(\frac{L'_\chi}{L_\chi} \right) (-a) \right] \\ &+ \frac{1}{x^a} \left[\left(\frac{\zeta'}{\zeta} \right)' (-a) - \left(\frac{L'_\chi}{L_\chi} \right)' (-a) \right]. \end{aligned}$$

Proof. Subtract Lemma 3.2 from Lemma 3.3. □

Lemma 3.13. *Let χ be a non-principal primitive Dirichlet character modulo f with $\chi(-1) = 1$. For $a \in (0, 1)$ we have*

$$\begin{aligned} &\left| \left(\frac{\zeta'}{\zeta} \right) (-a) - \left(\frac{L'_\chi}{L_\chi} \right) (-a) \right| \\ &\leq (a+2) \sum_{\rho \text{ of } \zeta, L_\chi} \frac{1}{|(\rho+a)(2-\rho)|} + 2 \left| \frac{\zeta'(2)}{\zeta(2)} \right| + \frac{1}{a} + \frac{1}{a+1} + \frac{3}{2}. \end{aligned}$$

Proof. We begin with the following formulas which hold for all $s \in \mathbb{C}$, provided the sums are taken in symmetric order:

$$(3.7) \quad \left(\frac{\zeta'}{\zeta} \right) (s) = \sum_{\rho \text{ of } \zeta} \frac{1}{s - \rho} - \frac{1}{s} - \frac{1}{s-1} - \psi_{\mathbb{Q}}(s)$$

$$(3.8) \quad \left(\frac{L'_\chi}{L_\chi} \right) (s) = \sum_{\rho \text{ of } L_\chi} \frac{1}{s - \rho} - \frac{1}{2} \log f - \psi_{\mathbb{Q}}(s)$$

Formulas (3.7) and (3.8) are obtained from (3.2) and (3.3) respectively by applying the facts $\sum_{\rho \text{ of } \zeta} \rho^{-1} + B = 0$ and $\sum_{\rho \text{ of } L_\chi} \rho^{-1} + B_\chi = 0$. Plugging $s = 2$ into (3.7) and subtracting it from itself, and similarly for (3.8), yields:

$$\begin{aligned} \left(\frac{\zeta'}{\zeta}\right)(s) &= \left(\frac{\zeta'}{\zeta}\right)(2) + \sum_{\rho} \left(\frac{1}{s-\rho} - \frac{1}{2-\rho}\right) \\ &\quad + \frac{3}{2} - \frac{1}{s} - \frac{1}{s-1} + \psi_{\mathbb{Q}}(2) - \psi_{\mathbb{Q}}(s) \\ \left(\frac{L'_\chi}{L_\chi}\right)(s) &= \left(\frac{L'_\chi}{L_\chi}\right)(2) + \sum_{\rho} \left(\frac{1}{s-\rho} - \frac{1}{2-\rho}\right) + \psi_{\mathbb{Q}}(2) - \psi_{\mathbb{Q}}(s) \end{aligned}$$

Using the above, together with the fact

$$\frac{1}{-a-\rho} - \frac{1}{2-\rho} = -\frac{a+2}{(\rho+a)(2-\rho)},$$

we can write

$$\begin{aligned} \left(\frac{\zeta'}{\zeta}\right)(-a) - \left(\frac{L'_\chi}{L_\chi}\right)(-a) &= (a+2) \left(\sum_{\rho \text{ of } L_\chi} \frac{1}{(\rho+a)(2-\rho)} - \sum_{\rho \text{ of } \zeta} \frac{1}{(\rho+a)(2-\rho)} \right) \\ &\quad + \left(\frac{\zeta'}{\zeta}\right)(2) - \left(\frac{L'_\chi}{L_\chi}\right)(2) + \frac{3}{2} + \frac{1}{a} + \frac{1}{a+1}. \end{aligned}$$

The result follows upon taking absolute values and using the fact that

$$\left| \left(\frac{L'_\chi}{L_\chi}\right)(2) \right| \leq \left| \left(\frac{\zeta'}{\zeta}\right)(2) \right|.$$

□

Lemma 3.14. *Suppose $a \in (0, 1)$ and $\Re(\rho) = 1/2$. Then*

$$\frac{1}{|(\rho+a)(2-\rho)|} \leq \frac{1}{|\rho+a|^2}.$$

Proof. Use $|2-\rho| \geq |\rho+a|$.

□

Lemma 3.15. *Let χ be a non-principal primitive Dirichlet character modulo f with $\chi(-1) = 1$. For $a \in (0, 1)$ we have*

$$\left| \left(\frac{\zeta'}{\zeta}\right)'(-a) - \left(\frac{L'_\chi}{L_\chi}\right)'(-a) \right| < \sum_{\rho \text{ of } \zeta, L_\chi} \frac{1}{|\rho+a|^2} + \frac{1}{a^2} + \frac{1}{(a+1)^2}.$$

Proof. We start by differentiating (3.7) and (3.8); this gives

$$(3.9) \quad \left(\frac{\zeta'}{\zeta}\right)'(s) = - \sum_{\rho \text{ of } \zeta} \frac{1}{(s-\rho)^2} + \frac{1}{s^2} + \frac{1}{(s-1)^2} - \psi'_{\mathbb{Q}}(s)$$

$$(3.10) \quad \left(\frac{L'_\chi}{L_\chi}\right)'(s) = - \sum_{\rho \text{ of } L_\chi} \frac{1}{(s-\rho)^2} - \psi'_{\mathbb{Q}}(s),$$

which allows us to write

$$\begin{aligned} & \left(\frac{\zeta'}{\zeta}\right)'(-a) - \left(\frac{L'_\chi}{L_\chi}\right)'(-a) \\ &= \sum_{\rho \text{ of } L_\chi} \frac{1}{(\rho+a)^2} - \sum_{\rho \text{ of } \zeta} \frac{1}{(\rho+a)^2} + \frac{1}{a^2} + \frac{1}{(a+1)^2}. \end{aligned}$$

The result follows. □

Proposition 3.1. *Let χ be a non-principal primitive Dirichlet character modulo f with $\chi(-1) = 1$. Assume the RH and the GRH for $L(s, \chi)$. We define*

$$\sum_{\rho} := \sum_{\rho \text{ of } \zeta, L_\chi} \frac{1}{\left|\rho + \frac{1}{2}\right|^2}.$$

For $x > 0$ we have

$$\begin{aligned} \frac{x}{9/4} + 4 &\leq \sqrt{x} \sum_{\rho} + 2 \sum_{\substack{n < x \\ \chi(n) \neq 1}} \Lambda(n)(n/x)^{1/2} \log(x/n) \\ &+ \frac{\log x}{\sqrt{x}} \left(\frac{5}{2} \sum_{\rho} + 2 \left| \frac{\zeta'(2)}{\zeta(2)} \right| + \frac{25}{6} \right) + \frac{1}{\sqrt{x}} \left(\sum_{\rho} + \frac{40}{9} \right). \end{aligned}$$

Proof. Set $a = 1/2$. Combine Lemmas 3.12, 3.13, 3.14, and 3.15. □

Proof of Theorem 3.2. The result for a general character follows from the corresponding result for primitive characters and hence we may assume χ is a primitive character modulo f .

Define $x := 2.5(\log f)^2$. Since $f \geq 10^9$, we have $x > 1073$. By way of contradiction, suppose that $\chi(n) = 1$ for all $n < x$ with $(n, q_1) = 1$. Under this assumption, we apply Lemma 3.4 with $u = q_1$, which gives

$$\sum_{\substack{n < x \\ \chi(n) \neq 1}} \Lambda(n)(n/x)^{1/2} \log(x/n) \leq (\log x)^2.$$

Combining the above with Proposition 3.1 and dividing by \sqrt{x} yields

$$\frac{\sqrt{x}}{9/4} + \frac{4}{\sqrt{x}} \leq \sum_{\rho} + \frac{2(\log x)^2}{\sqrt{x}} + \frac{\log x}{x} \left[\frac{5}{2} \sum_{\rho} + 2 \left| \frac{\zeta'(2)}{\zeta(2)} \right| + \frac{25}{6} \right] + \frac{1}{x} \left[\sum_{\rho} + \frac{40}{9} \right].$$

By Lemma 3.7, we have

$$\sum_{\rho} \leq \frac{1}{2} \log f + 0.437,$$

and in particular,

$$\frac{1}{\sqrt{x}} \sum_{\rho} \leq \frac{1}{3}.$$

We see

$$\frac{\log x}{x} \left[\frac{25}{6} \right] + \frac{1}{x} \left[\sum_{\rho} + \frac{40}{9} \right] \leq \frac{1}{\sqrt{x}} \left(\frac{\log x}{\sqrt{x}} \cdot \frac{25}{6} + \frac{1}{3} + \frac{1}{\sqrt{x}} \frac{40}{9} \right) < \frac{4}{\sqrt{x}},$$

and therefore

$$\frac{\sqrt{x}}{9/4} \leq \sum_{\rho} + \frac{2(\log x)^2}{\sqrt{x}} + \frac{\log x}{x} \left[\frac{5}{2} \sum_{\rho} + 2 \left| \frac{\zeta'(2)}{\zeta(2)} \right| \right].$$

We have

$$\frac{1}{\sqrt{x}} \left[\frac{5}{2} \sum_{\rho} + 2 \left| \frac{\zeta'(2)}{\zeta(2)} \right| \right] < 0.869$$

and

$$\frac{\log x}{\sqrt{x}} \leq 0.214$$

which leads to

$$\frac{\sqrt{x}}{9/4} \leq \frac{1}{2} \log f + 0.437 + \frac{2(\log x)^2}{\sqrt{x}} + 0.186.$$

Now we observe

$$\frac{2(\log x)^2}{\sqrt{x}} \leq 2.98.$$

All together, we have

$$\frac{\sqrt{x}}{9/4} \leq \frac{1}{2} \log f + 3.61$$

This leads to:

$$\begin{aligned} \sqrt{x} &\leq \frac{9}{8} (\log f) + 8.13 \\ &\leq 1.52 \log f \end{aligned}$$

Squaring both sides yields

$$x \leq 2.32 (\log f)^2,$$

a contradiction. □

3.4. An upper estimate on r . We establish a series of results, building up to the proof of Theorem 3.3.

Lemma 3.16. *Let χ be a non-principal Dirichlet character modulo a prime p of order ℓ with $\chi(-1) = 1$. Fix any ℓ -th root of unity $\omega \neq 1$. For $a \in (0, 1)$ and $x \in (1, p)$ we have*

$$\begin{aligned} \frac{x}{(a+1)^2} + \frac{1}{a^2} &= \sum_{k=1}^{\ell} \omega^{-k} \sum_{\rho \text{ of } L_{\chi^k}} \frac{x^\rho}{(\rho+a)^2} \\ &\quad + \ell \sum_{\substack{n < x \\ \chi(n) = \omega}} \Lambda(n) (n/x)^a \log(x/n) \\ &\quad + \frac{\log x}{x^a} \left[\left(\frac{\zeta'}{\zeta} \right) (-a) + \sum_{k=1}^{\ell-1} \omega^{-k} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right) (-a) \right] \\ &\quad + \frac{1}{x^a} \left[\left(\frac{\zeta'}{\zeta} \right)' (-a) + \sum_{k=1}^{\ell-1} \omega^{-k} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right)' (-a) \right]. \end{aligned}$$

Proof. First we note that χ^k for $k = 1, \dots, \ell - 1$ are all non-principal primitive characters as χ is a character modulo a prime p of order ℓ ; moreover, $\chi^\ell(n) = 1$ for all $n < x$ as $x < p$. Multiplying the identity

$$\sum_{k=1}^{\ell} \omega^{-k} \chi^k(n) = \begin{cases} \ell & \chi(n) = \omega \\ 0 & \text{otherwise} \end{cases}$$

by

$$g(x, n) := \Lambda(n) (n/x)^a \log(x/n)$$

and summing over all $n < x$ yields

$$\sum_{n < x} g(x, n) \sum_{k=1}^{\ell} \omega^{-k} \chi^k(n) = \ell \sum_{\substack{n < x \\ \chi(n) = \omega}} g(x, n).$$

Interchanging the order of summation gives

$$\sum_{k=1}^{\ell} \omega^{-k} \sum_{n < x} g(x, n) \chi^k(n) = \ell \sum_{\substack{n < x \\ \chi(n) = \omega}} g(x, n).$$

Now we apply Lemma 3.2 and Lemma 3.3 and use the facts:

$$(3.11) \quad \sum_{k=1}^{\ell} \omega^{-k} = 0, \quad \sum_{k=1}^{\ell-1} \omega^{-k} = -1.$$

The result follows. □

Lemma 3.17. *Let χ be a non-principal Dirichlet character modulo a prime p of order ℓ with $\chi(-1) = 1$. Fix any ℓ -th root of unity $\omega \neq 1$. For $a \in (0, 1)$ we have*

$$\begin{aligned} & \left| \left(\frac{\zeta'}{\zeta} \right) (-a) + \sum_{k=1}^{\ell-1} \omega^{-k} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right) (-a) \right| \\ & \leq (a+2) \sum_{\rho} \frac{1}{|(\rho+a)(2-\rho)|} + \ell \left| \frac{\zeta'(2)}{\zeta(2)} \right| + \frac{1}{a} + \frac{1}{a+1} + \frac{3}{2}, \end{aligned}$$

where the sum is taken over all non-trivial zeros ρ of $L(s, \chi^k)$ for $k = 1, \dots, \ell$.

Proof. Using (3.7), (3.8) and (3.11), we can write:

$$\begin{aligned} & \left(\frac{\zeta'}{\zeta} \right) (-a) + \sum_{k=1}^{\ell-1} \omega^{-k} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right) (-a) \\ & = -(a+2) \sum_{k=1}^{\ell} \omega^{-k} \sum_{\rho \text{ of } L_{\chi^k}} \frac{1}{(\rho+a)(2-\rho)} \\ & \quad + \left(\frac{\zeta'}{\zeta} \right) (2) + \sum_{k=1}^{\ell-1} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right) (2) + \frac{3}{2} + \frac{1}{a} + \frac{1}{a+1} \end{aligned}$$

The result follows in a similar manner as Lemma 3.13. □

Lemma 3.18. *Let χ be a non-principal Dirichlet character modulo a prime p of order ℓ with $\chi(-1) = 1$. Fix any ℓ -th root of unity $\omega \neq 1$. For $a \in (0, 1)$ we have*

$$\left| \left(\frac{\zeta'}{\zeta} \right)' (-a) + \sum_{k=1}^{\ell-1} \omega^{-k} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right)' (-a) \right| \leq \sum_{\rho} \frac{1}{|\rho+a|^2} + \frac{1}{a^2} + \frac{1}{(a+1)^2},$$

where the sum is taken over all non-trivial zeros ρ of $L(s, \chi^k)$ for $k = 1, \dots, \ell$.

Proof. Using (3.9), (3.10), and (3.11) we can write

$$\begin{aligned} & \left(\frac{\zeta'}{\zeta} \right)' (-a) + \sum_{k=1}^{\ell-1} \omega^{-k} \left(\frac{L'_{\chi^k}}{L_{\chi^k}} \right)' (-a) \\ & = \frac{1}{a^2} + \frac{1}{(a+1)^2} - \sum_{k=1}^{\ell} \omega^{-k} \sum_{\rho \text{ of } L_{\chi^k}} \frac{1}{(\rho+a)^2}. \end{aligned}$$

□

Proposition 3.2. *Let ℓ and f be odd primes with $f \equiv 1 \pmod{\ell}$. Let K be the Galois number field of degree ℓ and conductor f , and let χ be a primitive Dirichlet character modulo f of order ℓ . Fix any ℓ -th root of unity $\omega \neq 1$. Suppose that the GRH holds for $\zeta_K(s)$. We define*

$$\sum_{\rho} := \sum_{\rho} \frac{1}{\left| \rho + \frac{1}{2} \right|^2},$$

where the sum is taken over all non-trivial zeros of $\zeta_K(s)$. For $x \in (1, f)$ we have

$$\begin{aligned} \frac{x}{9/4} + 4 \leq \sqrt{x} \sum_{\rho} + \ell \sum_{\substack{n < x \\ \chi(n) = \omega}} \Lambda(n)(n/x)^{1/2} \log(x/n) \\ + \frac{\log x}{\sqrt{x}} \left(\frac{5}{2} \sum_{\rho} + \ell \left| \frac{\zeta'(2)}{\zeta(2)} \right| + \frac{25}{6} \right) + \frac{1}{\sqrt{x}} \left(\sum_{\rho} + \frac{40}{9} \right). \end{aligned}$$

Proof. In light of Lemma 2.1, \sum_{ρ} can also be thought of as the sum over the non-trivial zeros of $L(s, \chi^k)$ for $k = 1, \dots, \ell$ (counting multiplicities). Observe that since ℓ is odd, we have $\chi(-1) = 1$. Now set $a = 1/2$ and combine Lemmas 3.16, 3.17, 3.14, 3.18. \square

Proof of Theorem 3.3. Define $x := 2.5(\ell - 1)^2(\log f)^2$. Since $f \geq 10^8$ and $\ell \geq 3$, we have $x > 3393$. By way of contradiction, suppose that $\chi(n) \neq \omega$ for all $n < x$ with $(n, q_1 q_2) = 1$. Under this assumption we apply Lemma 3.4 with $u = q_1 q_2$, which gives

$$\sum_{\substack{n < x \\ \chi(n) = \omega}} \Lambda(n)(n/x)^{1/2} \log(x/n) \leq 2(\log x)^2.$$

Combining the above with Proposition 3.2 and dividing by \sqrt{x} yields:

$$\frac{\sqrt{x}}{9/4} + \frac{4}{\sqrt{x}} \leq \sum_{\rho} + \frac{2\ell(\log x)^2}{\sqrt{x}} + \frac{\log x}{x} \left[\frac{5}{2} \sum_{\rho} + \ell \left| \frac{\zeta'(2)}{\zeta(2)} \right| + \frac{25}{6} \right] + \frac{1}{x} \left[\sum_{\rho} + \frac{40}{9} \right].$$

We note that Lemma 3.11 is applicable in our situation; indeed, our assumptions on K imply that it is totally-real and, using the conductor-discriminant formula, we see that $\Delta = f^{\ell-1}$. By Lemma 3.11, we have

$$\sum_{\rho} \leq \frac{1}{2} [(\ell - 1) \log f - 2.23 \ell + 5.34],$$

and in particular,

$$\frac{1}{\sqrt{x}} \sum_{\rho} \leq \frac{1}{2\sqrt{2.5}}.$$

We see

$$\frac{\log x}{x} \left[\frac{25}{6} \right] + \frac{1}{x} \left[\sum_{\rho} + \frac{40}{9} \right] \leq \frac{1}{\sqrt{x}} \left(\frac{\log x}{\sqrt{x}} \cdot \frac{25}{6} + \frac{1}{2\sqrt{2.5}} + \frac{1}{\sqrt{x}} \frac{40}{9} \right) < \frac{4}{\sqrt{x}},$$

and therefore

$$\frac{\sqrt{x}}{9/4} \leq \sum_{\rho} + \frac{2\ell(\log x)^2}{\sqrt{x}} + \frac{\log x}{x} \left[\frac{5}{2} \sum_{\rho} + \ell \left| \frac{\zeta'(2)}{\zeta(2)} \right| \right].$$

We have

$$\frac{1}{\sqrt{x}} \left[\frac{5}{2} \sum_{\rho} + \ell \left| \frac{\zeta'(2)}{\zeta(2)} \right| \right] < 0.82$$

and

$$\frac{\log x}{\sqrt{x}} \leq 0.14,$$

which leads to

$$\frac{\sqrt{x}}{9/4} \leq \frac{1}{2} [(\ell - 1) \log f - 2.23 \ell + 5.34] + \frac{2\ell(\log x)^2}{\sqrt{x}} + 0.12.$$

Now we observe

$$\frac{2\ell(\log x)^2}{\sqrt{x}} \leq 2.27\ell.$$

All together, we have

$$\begin{aligned} \frac{\sqrt{x}}{9/4} &\leq \frac{1}{2} [(\ell - 1) \log f - 2.23 \ell + 5.34] + 2.27\ell + 0.12 \\ &\leq \frac{1}{2} (\ell - 1) (\log f) + 1.16\ell + 2.79 \end{aligned}$$

This leads to:

$$\begin{aligned} \sqrt{x} &\leq \frac{9}{8} (\ell - 1) (\log f) + 2.61\ell + 6.28 \\ &\leq 1.51(\ell - 1) \log f \end{aligned}$$

Squaring both sides yields

$$x \leq 2.3(\ell - 1)^2 (\log f)^2,$$

a contradiction. □

4. GRH bounds for norm-Euclidean fields

In this section we prove Theorem 1.4. First we deal separately with the situation where q_1 is small.

Theorem 4.1. *Let K be a Galois number field of odd prime degree ℓ and conductor f . Assume the GRH for $\zeta_K(s)$. Let q_1 denote the smallest rational prime which is inert in K . If $q_1 < 100$ and*

$$5825(\ell - 1)^2(\log f)^4 < f,$$

then K is not norm-Euclidean.

Proof. Set $A = 5825$. One checks that our hypothesis implies $f \geq 10^9$. By Theorem 2.1 we may assume that f is a prime with $f \equiv 1 \pmod{\ell}$. We adopt the notation from the statement of Theorem 2.2. Since $q_1 < 100$, we have $q_1 \leq 97$, and by Theorems 3.2 and 3.3, we have

$$(4.1) \quad q_2 < 2.5(\log f)^2,$$

$$(4.2) \quad r < 2.5(\ell - 1)^2(\log f)^2;$$

hence we have

$$\begin{aligned} 2.1 q_1 q_2 r \log q_1 &< (2.1)(97)(2.5)(\log f)^2(2.5)(\ell - 1)^2(\log f)^2(\log 97) \\ &< A(\ell - 1)^2(\log f)^4. \end{aligned}$$

If $q_1 \neq 2, 3, 7$, then it follows from Theorem 2.2 that the condition given in our hypothesis is sufficient. If $q_1 = 7$, then we observe that

$$\begin{aligned} 3 q_1 q_2 r \log q_1 &< (3)(7)(2.5)(\log f)^2(2.5)(\ell - 1)^2(\log f)^2(\log 7) \\ &< A(\ell - 1)^2(\log f)^4. \end{aligned}$$

Now we deal with the special case where $q_1 = 2$, $q_2 = 3$. Our hypothesis gives

$$(\ell - 1) < \frac{1}{\sqrt{A}} \frac{f^{1/2}}{(\log f)^2}.$$

In order to use Proposition 2.1, we estimate

$$\begin{aligned} 72(\ell - 1)f^{1/2} \log 4f + 35 &< \frac{72}{\sqrt{A}} \frac{\log 4f}{(\log f)^2} f + 35 \\ &< 0.1f + 35 \\ &< f; \end{aligned}$$

thus the proposition applies. When $q_1 = 3$, $q_2 = 5$, we use a similar estimate to conclude that

$$507(\ell - 1)f^{1/2} \log 9f + 448 < 0.4f + 448 < f,$$

and hence Proposition 2.1 applies again.

The remaining cases fall under conditions (4) and (5) of Theorem 2.2. We will prove the bound

$$5 q_2 r < f,$$

which will deal with all remaining cases. From the estimates (4.1) and (4.2) we have

$$5 q_2 r < 32(\ell - 1)^2(\log f)^4 < A(\ell - 1)^2(\log f)^4 < f.$$

This completes the proof. □

Applying the previous theorem with $\ell = 3$ yields:

Corollary 4.1. *Let K be a Galois cubic number field with conductor $f \geq 6 \cdot 10^9$. Assume the GRH for $\zeta_K(s)$. Let q_1 denote the smallest rational prime which is inert in K . If $q_1 < 100$, then K is not norm-Euclidean.*

Proof of Theorem 1.4. One checks that our hypothesis implies $f \geq 10^{10}$. By Theorem 2.1 we may assume that f is a prime with $f \equiv 1 \pmod{\ell}$. We adopt the notation from Theorem 2.2. Applying Theorems 3.1, 3.2, and 3.3, we have:

$$(4.3) \quad \begin{aligned} q_1 &< (1.17 \log f - 6.3)^2 \\ q_2 &\leq 2.5(\log f)^2 \end{aligned}$$

$$(4.4) \quad r \leq 2.5(\ell - 1)^2(\log f)^2$$

For the moment, we assume $q_1 \neq 2, 3, 7$. Combining everything, this gives $2.1 q_1 q_2 r \log q_1 < 26.25(\ell - 1)^2(1.17 \log f - 6.3)^2 \log(1.17 \log f - 6.3)(\log f)^4$. Hence a sufficient condition is:

$$(4.5) \quad 26.25(\ell - 1)^2(1.17 \log f - 6.3)^2 \log(1.17 \log f - 6.3)(\log f)^4 \leq f$$

Note that the condition given in our hypothesis implies (4.5). To deal with the remaining cases of $q_1 = 2, 3, 7$, we note that (4.5) implies the condition given in the statement of Theorem 4.1; hence (4.5) is sufficient in all cases. □

5. Galois cubic fields

Finally, we give the proof of Theorem 1.1. Let K be a norm-Euclidean Galois cubic field with conductor f and discriminant Δ which is not any of the 13 fields listed in the statement of Theorem 1.1. In light of Theorem 1.2, we may assume $f \geq 10^{10}$. Moreover, Theorem 2.1 allows us to conclude that $\Delta = f^2$ and that f is a prime with $f \equiv 1 \pmod{3}$. Using the slightly complicated condition (4.5) in the proof of Theorem 1.4 and setting $\ell = 3$ we find that $f < 7 \cdot 10^{10}$.

It remains to deal with the cases where f lies in $(10^{10}, 7 \cdot 10^{10})$. Let χ be a primitive cubic character modulo f , and let q_1 denote the smallest

prime such that $\chi(q_1) \neq 1$. By Corollary 4.1, to show that K is not norm-Euclidean, assuming $f \in (10^{10}, 7 \cdot 10^{10})$, it suffices to show $q_1 < 100$. Using a computer program (written in C) to carry out the computation, we obtain the following lemma which completes the proof of Theorem 1.1.

Lemma 5.1. *Suppose f is a prime with $f \equiv 1 \pmod{3}$. Let χ be a cubic character modulo f , and denote by q_1 the smallest prime with $\chi(q_1) \neq 1$. If $f \leq 7 \cdot 10^{10}$, then $q_1 \leq 61$.*

In our implementation we use NTL with GMP for large integer arithmetic. We use the fact that $\chi(p) = 1$ if and only if $p^{(f-1)/3} \equiv 1 \pmod{f}$, as the latter condition can be checked very quickly using fast modular exponentiation. The computation was carried out on a MacBook Pro with a 2.26 GHz Intel Core 2 Duo processor and 4 GB of RAM, running Mac OS 10.6. It took approximately 4 hours of CPU time to complete.

As an additional curiosity we have kept a list of record values of q_1 . That is, each time we encounter a value of q_1 which is strictly greater than all previous values, we have outputted the values of f and q_1 . Here are the results:

```
Record: f=7, q1=2
Record: f=31, q1=3
Record: f=307, q1=5
Record: f=643, q1=7
Record: f=5113, q1=11
Record: f=21787, q1=13
Record: f=39199, q1=17
Record: f=360007, q1=23
Record: f=4775569, q1=29
Record: f=10318249, q1=37
Record: f=65139031, q1=41
Record: f=387453811, q1=43
Record: f=913900417, q1=47
Record: f=2278522747, q1=53
Record: f=2741702809, q1=59
Record: f=25147657981, q1=61
```

Acknowledgement

This paper is based upon a portion of the author's Ph.D. dissertation under the supervision of Professor Harold Stark. The author would like to thank Professor Stark for his invaluable guidance and support at all stages of this work.

References

- [1] N. C. ANKENY, *The least quadratic non residue*. Ann. of Math. (2) **55** (1952), 65–72.
- [2] E. BACH, *Explicit bounds for primality testing and related problems*. Math. Comp. **55** (1990), no. 191, 355–380.

- [3] H. DAVENPORT, *Multiplicative number theory, third edition*. Graduate Texts in Mathematics, vol. **74**, Springer-Verlag, New York, 2000, Revised and with a preface by Hugh L. Montgomery.
- [4] H. J. GODWIN AND J. R. SMITH, *On the Euclidean nature of four cyclic cubic fields*. Math. Comp. **60** (1993), no. 201, 421–423.
- [5] H. HEILBRONN, *On Euclid’s algorithm in cubic self-conjugate fields*. Proc. Cambridge Philos. Soc. **46** (1950), 377–382.
- [6] Y. IHARA, V. K. MURTY, AND M. SHIMURA, *On the logarithmic derivatives of Dirichlet L -functions at $s = 1$* . Acta Arith. **137** (2009), no. 3, 253–276.
- [7] J. C. LAGARIAS AND A. M. ODLYZKO, *Effective versions of the Chebotarev density theorem*. Algebraic number fields: L -functions and Galois properties (Proc. Sympos., Univ. Durham, Durham, 1975), Academic Press, London, 1977, pp. 409–464.
- [8] E. LANDAU, *Zur Theorie der Heckschen Zetafunktionen, welche komplexen Charakteren entsprechen*. Math. Z. **4** (1919), no. 1-2, 152–162.
- [9] F. LEMMERMEYER, *The Euclidean algorithm in algebraic number fields*. Exposition. Math. **13** (1995), no. 5, 385–416.
- [10] K. J. MCGOWN, *Norm-Euclidean cyclic fields of prime degree*. Int. J. Number Theory (to appear).
- [11] H. L. MONTGOMERY, *Topics in multiplicative number theory*. Lecture Notes in Mathematics, Vol. **227**, Springer-Verlag, Berlin, 1971.
- [12] W. NARKIEWICZ, *Elementary and analytic theory of algebraic numbers, second edition*. Springer-Verlag, Berlin, 1990.
- [13] A. M. ODLYZKO, *Bounds for discriminants and related estimates for class numbers, regulators and zeros of zeta functions: a survey of recent results*. Sémin. Théor. Nombres Bordeaux (2) **2** (1990), no. 1, 119–141.
- [14] G. POITOU, *Minorations de discriminants (d’après A. M. Odlyzko)*. Séminaire Bourbaki, Vol. 1975/76 28ème année, Exp. No. 479, Springer, Berlin, 1977, pp. 136–153. Lecture Notes in Math., Vol. **567**.
- [15] G. POITOU, *Sur les petits discriminants*. Séminaire Delange-Pisot-Poitou, 18e année: (1976/77), Théorie des nombres, Fasc. 1, Secrétariat Math., Paris, 1977, pp. Exp. No. 6, 18.
- [16] H. M. STARK, *Some effective cases of the Brauer-Siegel theorem*. Invent. Math. **23** (1974), 135–152.
- [17] H. M. STARK, *The analytic theory of algebraic numbers*. Bull. Amer. Math. Soc. **81** (1975), no. 6, 961–972.
- [18] A. WEIL, *Sur les “formules explicites” de la théorie des nombres premiers*. Comm. Sémin. Math. Univ. Lund (1952), 252–265.
- [19] A. WEIL, *Sur les formules explicites de la théorie des nombres*. Izv. Akad. Nauk SSSR Ser. Mat. **36** (1972), 3–18.

Kevin J. MCGOWN
 Department of Mathematics
 University of California, San Diego
 La Jolla, California, 92093, USA

Current address :

Department of Mathematics
 Oregon State University
 368 Kidder Hall
 Corvallis, Oregon, 97331, USA
E-mail: mcgownk@math.oregonstate.edu