

# JOURNAL

de Théorie des Nombres  
de BORDEAUX

*anciennement Séminaire de Théorie des Nombres de Bordeaux*

Paul FILI

**On the heights of totally  $p$ -adic numbers**

Tome 26, n° 1 (2014), p. 103-109.

<[http://jtnb.cedram.org/item?id=JTNB\\_2014\\_\\_26\\_1\\_103\\_0](http://jtnb.cedram.org/item?id=JTNB_2014__26_1_103_0)>

© Société Arithmétique de Bordeaux, 2014, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du*  
*Centre de diffusion des revues académiques de mathématiques*  
<http://www.cedram.org/>

## On the heights of totally $p$ -adic numbers

par PAUL FILI

RÉSUMÉ. Bombieri et Zannier ont démontré des minoration et des majorations de la limite inférieure de la hauteur de Weil sur le corps des nombres totalement  $p$ -adiques et sur leurs généralisations. Dans notre étude nous utilisons des techniques de la théorie du potentiel pour généraliser les majorations de leur étude et, dans l'hypothèse d'intégralité, améliorer un peu plus les minoration.

ABSTRACT. Bombieri and Zannier established lower and upper bounds for the limit infimum of the Weil height in fields of totally  $p$ -adic numbers and generalizations thereof. In this paper, we use potential theoretic techniques to generalize the upper bounds from their paper and, under the assumption of integrality, to improve slightly upon their bounds.

### 1. Statement of Results

Recall that an algebraic number is said to be *totally  $p$ -adic* if its image lies in  $\mathbb{Q}_p$  for any embedding  $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ , where  $\mathbb{C}_p$  denotes the completion of an algebraic closure of  $\mathbb{Q}_p$ . This is analogous to the usual definition of a totally real number, however, unlike  $\mathbb{C}/\mathbb{R}$ , the extension  $\mathbb{C}_p/\mathbb{Q}_p$  is of infinite degree, so in fact we can make an even broader generalization:

**Definition 1.** Let  $L_p/\mathbb{Q}_p$  be a (finite) Galois extension for  $p \leq \infty$  a rational prime. We say  $\alpha \in \overline{\mathbb{Q}}$  is *totally  $L_p$*  if all Galois conjugates of  $\alpha$  lie in  $L_p \subset \mathbb{C}_p$ .

More generally, when our objects are defined over an arbitrary number field  $K$ , we make the following definition:

**Definition 2.** Fix a base number field  $K$ , and let  $S$  be a set of places of  $K$ . For each  $v \in S$ , we choose a Galois extension  $L_v/K_v$ . We say that  $\alpha$  is *totally  $L_S/K$*  if, for each  $v \in S$ , all of the  $K$ -Galois conjugates of  $\alpha$  lie in  $L_v$ .

Notice that  $\alpha$  is totally  $L_S/K$  if and only if the minimal polynomial for  $\alpha$  over  $K$  splits in  $L_v$  for each  $v \in S$ . In our terminology, a number being totally real is equivalent to being totally  $\mathbb{R}$  (or totally  $\mathbb{R}/\mathbb{Q}$ ), and being

totally  $p$ -adic is equivalent to being totally  $\mathbb{Q}_p$  (totally  $\mathbb{Q}_p/\mathbb{Q}$ ). Notice that the set of all totally  $L_S/K$  numbers form a normal extension of  $K$  (typically of infinite degree).

Bombieri and Zannier [3] studied the question of what the limit infimum of the Weil height was in these fields when the base field was assumed to be  $\mathbb{Q}$ , and they proved the following:

**Theorem** (Bombieri and Zannier 2001). *Let  $S$  is the set of finite rational primes and for each  $p \in S$ , let  $L_p/\mathbb{Q}_p$  be a chosen Galois extension. Suppose  $L/\mathbb{Q}$  is a normal extension (possibly of infinite degree) such that  $L$  embeds in  $L_p$  for each  $p \in S$ . Then*

$$\liminf_{\alpha \in L} h(\alpha) \geq \frac{1}{2} \sum_{p \in S} \frac{\log p}{e_p(p^{f_p} + 1)}$$

where  $e_p$  and  $f_p$  denote the ramification and inertial degrees of  $L_p/\mathbb{Q}_p$ , respectively.

Further, in the case where  $S = \{p_1, \dots, p_n\}$  is a finite set of nonarchimedean rational primes with  $L_{p_i} = \mathbb{Q}_{p_i}$  for each  $i$ , if we let  $L$  be the field of all totally  $L_S/\mathbb{Q}$  numbers, then we have

$$\liminf_{\alpha \in L} h(\alpha) \leq \sum_{i=1}^n \frac{\log p_i}{p_i - 1}.$$

We note that in Bombieri and Zannier's result, by the assumption that our extensions are normal, it follows that every  $p$ -adic completion of  $L$  has ramification and inertial degree less than or equal to that of  $L_p$ . More simply, we might take  $L$  to be the field of all totally  $L_S/\mathbb{Q}$  numbers above and recover the same result.

Smyth [8, 9] and Flammang [4] proved analogous results in the totally real case for the limit infimum of the height, however, they imposed the additional hypothesis that one consider only totally real algebraic integers.

This note has two aims: first, to prove a slightly sharper lower bound by adding the hypothesis that we consider only algebraic integers, and second, to generalize the upper bound from Bombieri and Zannier's paper by the aid of the Fekete-Szegő theorem with splitting conditions, as formulated and proven by Rumely [5, 6]. Our results are the following:

**Theorem 1.1.** *Fix a number field  $K$ , a set of nonarchimedean places  $S$  of  $K$ , and a choice of Galois extension  $L_v/K_v$  for each  $v \in S$ . Let  $L$  be the field of all totally  $L_S/K$  numbers, and  $O_L$  denote its ring of integers. Then*

$$\liminf_{\alpha \in O_L} h(\alpha) \geq \frac{1}{2} \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}$$

where  $N_v = [K_v : \mathbb{Q}_v]/[K : \mathbb{Q}]$ ,  $e_v$  and  $f_v$  denote the ramification and inertial degrees of  $L_v/K_v$ , respectively, and  $q_v$  denotes the order of the residue field of  $K_v$ , and  $p_v$  is rational prime above which  $v$  lies.

As Bombieri and Zannier remark in their note, if the above sum diverges (which may happen if we allow  $S$  to be infinite), then it follows that the set of totally  $L_S/K$  algebraic integers satisfies the Northcott property (there are only a finite number with height below any fixed constant).

If we restrict our attention to extensions normal over  $\mathbb{Q}$ , as Bombieri and Zannier do, our result reduces to:

**Corollary 1.** *Fix a set of finite rational primes  $S$  and a choice of Galois extension  $L_p/\mathbb{Q}_p$  for each  $p \in S$ . Let  $L$  be the field of all totally  $L_S$  numbers, and  $O_L$  denote its ring of integers. Then*

$$\liminf_{\alpha \in O_L} h(\alpha) \geq \frac{1}{2} \sum_{p \in S} \frac{\log p}{e_p(p^{f_p} - 1)}$$

where  $e_p$  and  $f_p$  denote the ramification and inertial degrees of  $L_p/\mathbb{Q}_p$ , respectively.

**Theorem 1.2.** *Fix a number field  $K$ , a finite set of nonarchimedean places  $S$  of  $K$ , and a choice of Galois extension  $L_v/K_v$  for each  $v \in S$  with ramification degree  $e_v$ , inertial degree  $f_v$ , and residue field degree  $q_v$  and characteristic  $p_v$ . Let  $L$  be the field of all totally  $L_S/K$  numbers. Then*

$$\liminf_{\alpha \in L} h(\alpha) \leq \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}$$

where  $N_v = [K_v : \mathbb{Q}_v]/[K : \mathbb{Q}]$ .

Notice that this differs from our lower bound only by the factor of  $1/2$ .

One interesting thing to note in Theorem 1.1 is that the shape of our bound comes directly from the formula for the logarithmic capacity of the ring of integers of  $L_v$  at each place. This connection to potential theory gives an interesting explanation for the shape of the observed bound, and suggests that by determining minimal energy measures on  $L_v$  one might further improve the bounds above. In fact, this connection inspires us to conjecture that in fact the upper bound in Theorem 1.2 is sharp:

**Conjecture 1.** *Fix a number field  $K$ , a finite set of nonarchimedean places  $S$  of  $K$ , and a choice of Galois extension  $L_v/K_v$  for each  $v \in S$  with ramification degree  $e_v$ , inertial degree  $f_v$ , and residue field of degree  $q_v$  and characteristic  $p_v$ . Let  $L$  be the field of all totally  $L_S/K$  numbers and  $O_L$  its ring of integers. Then*

$$\liminf_{\alpha \in O_L} h(\alpha) = \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}$$

where  $N_v = [K_v : \mathbb{Q}_v]/[K : \mathbb{Q}]$ .

It is still an interesting open question whether the limit infimum for all numbers can be in fact achieved with algebraic integers or not. In this direction, we will note only that the result of Bombieri and Zannier, when read as an equidistribution result, seems to indicate that totally  $p$ -adic points of low height should be distributing evenly in the residue classes of  $\mathbb{P}^1(\mathbb{F}_p)$ . This seems to suggest that the limit infimum over an entire field  $L$  of the type constructed above might be smaller than that obtained by algebraic integers; however, all of the smallest known limit points, both for the height of totally real numbers (see [8]) and the height of totally  $p$ -adic numbers (as in Theorem 1.2) are in fact achieved by sequences of algebraic integers.

## 2. Proofs

*Proof of Theorem 1.1.* We now proceed to prove Theorem 1.1. Recall that we have fixed a base number field  $K$ , and let  $M_K$  denotes the places of  $K$ . First, let us note that if  $S$  is infinite, we can take a limit over increasing finite subsets of  $S$ , and the general result will follow, so we may as well assume that  $S$  is finite in our proof. For convenience let

$$N_v = \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]}.$$

We recall M. Baker's reformulation of Mahler's inequality from [1], namely, if  $\alpha$  is an algebraic number and  $\alpha_1, \dots, \alpha_n$  denote its  $K$ -Galois conjugates, then

$$(2.1) \quad \frac{1}{n(n-1)} \sum_{i \neq j} g_v(x, y) \geq \begin{cases} -N_v \cdot \frac{\log n}{n-1} & \text{if } v \mid \infty, \\ 0 & \text{otherwise} \end{cases}$$

where

$$g_p(x, y) = \log^+ |x|_v + \log^+ |y|_v - \log |x - y|_v,$$

and our absolute values are normalized so that  $|\cdot|_v = \|\cdot\|_v^{N_v}$  where  $\|\cdot\|_v$  extends the usual absolute of  $\mathbb{Q}$  over which it lies (thus the absolute values  $|\cdot|_v$  satisfy the product formula, and  $h(\alpha) = \sum_v \log^+ |\alpha|_v$ ). We assume in our theorem that  $\alpha$  is integral. The key role is played by the discriminant here. At the place  $v \in M_K$  ( $p$  here being the rational prime over which the place  $v$  lies), all of the conjugates of  $\alpha$  lie in ring of integers  $O_{L_v}$ . By [7, Example 4.1.24], we have that the  $v$ -adic logarithmic capacity of  $O_{L_v}$  with respect to the point  $\infty$ , which we will denote  $\gamma_{\infty, v}(O_{L_v})$ , satisfies

$$\log \gamma_{\infty, v}(O_{L_v}) = -N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)} < 0,$$

where  $q_v$  denotes the order of the residue field of  $O_{L_v}$  and  $p_v$  its characteristic.<sup>1</sup> We can also view this as the transfinite diameter of  $O_{L_v}$ ; more specifically, if we adopt the notation of [2, Ch. 6] and let

$$\log d_n(O_{L_v})_\infty = \sup_{z_1, \dots, z_n \in O_{L_v}} \frac{1}{n(n-1)} \log |z_i - z_j|_v,$$

then by [2, Lemma 6.21 and Theorem 6.23] it follows that<sup>2</sup>

$$\lim_{n \rightarrow \infty} \log d_n(O_{L_v})_\infty = \log \gamma_{\infty, v}(O_{L_v}).$$

Let  $\{\alpha^{(k)}\}_{k=1}^\infty$  denote a sequence of algebraic numbers such that

$$\lim_{k \rightarrow \infty} h(\alpha^{(k)}) = \liminf_{\alpha \in O_L} h(\alpha),$$

and let  $n_k$  denote the number of  $K$ -Galois conjugates of  $\alpha^{(k)}$ , which we denote  $\alpha_1^{(k)}, \dots, \alpha_{n_k}^{(k)}$ . Since the  $\alpha^{(k)}$  have bounded height, it follows from Northcott's theorem that  $n_k \rightarrow \infty$  as  $k \rightarrow \infty$ . By definition of  $d_n(O_{L_v})_\infty$ , we have

$$\frac{1}{n_k(n_k-1)} \sum_{i \neq j} \log |\alpha_i^{(k)} - \alpha_j^{(k)}|_v \leq \log d_{n_k}(O_{L_v})_\infty$$

for any totally  $L_S/K$  integer and  $v \in S$ , so it follows that

$$\limsup_{k \rightarrow \infty} \sum_{v \in S} \frac{1}{n_k(n_k-1)} \sum_{i \neq j} \log |\alpha_i^{(k)} - \alpha_j^{(k)}|_v \leq \sum_{v \in S} \log \gamma_{\infty, v}(O_{L_v}).$$

Therefore, by the product formula, using the assumption of integrality,

$$\liminf_{k \rightarrow \infty} \sum_{\substack{w \in M_K \\ w|_\infty}} \frac{1}{n_k(n_k-1)} \sum_{i \neq j} \log |\alpha_i^{(k)} - \alpha_j^{(k)}|_w \geq - \sum_{v \in S} \log \gamma_{\infty, v}(O_{L_v}).$$

It now follows from (2.1) that

$$\begin{aligned} \liminf_{k \rightarrow \infty} 2h(\alpha^{(k)}) &\geq \liminf_{k \rightarrow \infty} \sum_{w|_\infty} \log^+ |\alpha^{(k)}|_w \\ &\geq - \sum_{v \in S} \log \gamma_{\infty, v}(O_{L_v}) = \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)}. \end{aligned}$$

Upon dividing each side by 2, the result follows. □

---

<sup>1</sup>In the text [7] the  $v$ -adic absolute value  $|\cdot|_v$  which is used to compute the capacity is normalized to agree with the modulus with respect to the additive Haar measure, so in the text a  $\log q_v$  appears in the numerator instead of  $\log p_v$ . We also wish to draw the reader's attention to the fact that the ring of integers  $O_{L_v}$  is 'strictly smaller' than the closed unit disc in  $\mathbb{P}^1(\mathbb{C}_v)$ , which has logarithmic capacity  $\log \gamma_{\infty, v}(D_v(0, 1)) = 0$ .

<sup>2</sup>Note that here for convenience our absolute values are normalized so that we have the same normalization factors as in computing the absolute logarithmic Weil height.

*Proof of Theorem 1.2.* We will proceed by applying the adelic Fekete-Szegő theorem with splitting conditions [6, Theorem 2.1] (see also [5] as well as [2, Theorem 6.27] for an interpretation in terms of the Berkovich topology). For each  $v \in S$ , let  $O_{L_v} \subset L_v$  denote the ring of integers in  $L_v$ . Notice that  $O_{L_v} \subset \mathbb{P}^1(\mathbb{C}_p)$  is a compact set and let  $\mathcal{D}_v(0, 1) \subset \mathbb{P}^1(\mathbb{C}_v)$  be the usual Berkovich unit disc. Let  $\mathbb{E} \subset \prod_{v \in M_K} \mathbb{A}^1(\mathbb{C}_v)$  be the adelic Berkovich set given by

$$\mathbb{E} = \prod_{v \in M_K} E_v \quad \text{where} \quad E_v = \begin{cases} O_{L_v} & \text{for } v \in S \\ \mathcal{D}_v(0, 1) & \text{for } v \notin S, \\ \mathcal{D}\left(0, \prod_{v \in S} q_v^{N_v/e_v(q_v^{f_v}-1)}\right) & \text{for } v \mid \infty. \end{cases}$$

By [7, Example 4.1.24] we have normalized  $v$ -adic logarithmic capacity

$$\log \gamma_{\infty, v}(O_{L_v}) = -N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)} \quad \text{for each } v \in S,$$

where  $\gamma_{\infty}$  denotes the capacity relative to  $\mathcal{X} = \{\infty\}$  in the notation of [6]. At the infinite places, we have

$$\sum_{w \mid \infty} \log \gamma_{\infty, w}(\mathcal{D}(0, \prod_{v \in S} q_v^{N_v/e_v(q_v^{f_v}-1)})) = \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)},$$

so that the adelic capacity, that is, the product of all of the normalized capacities, is

$$\gamma_{\infty}(\mathbb{E}) = \prod_{v \in M_K} \gamma_{\infty, v}(E_v) = 1.$$

Further, by our assumptions,  $\mathbb{E}$  is stable under the action of the continuous Galois automorphisms of  $\mathbb{C}_v/K_v$  at each place. For  $\epsilon > 0$ , we consider the Berkovich adelic neighborhood

$$\mathbb{U} = \prod_{w \in M_K} U_w \quad \text{where} \quad U_w = \begin{cases} E_w & \text{for } w \nmid \infty \\ \mathcal{D}\left(0, e^{\epsilon} \prod_{v \in S} q_v^{N_v/e_v(q_v^{f_v}-1)}\right)^{-} & \text{for } w \mid \infty, \end{cases}$$

where  $\mathcal{D}(a, r)^{-}$  denotes the usual open disc centered at  $a$  of radius  $r$ . Then we can apply the adelic Fekete-Szegő theorem with splitting conditions as formulated in [6, Theorem 2.1], there are infinitely many algebraic numbers in  $\overline{K}$  all of whose conjugates lie in  $\mathbb{U}$  with the additional condition that all  $\mathbb{C}_v/K_v$  conjugates lie in  $U_v = E_v = O_{L_v}$  for each  $v \in S$ . Clearly, the only contribution to the height of such numbers comes from the archimedean places, and as a result, we have

$$h(\alpha) \leq \sum_{v \in S} N_v \cdot \frac{\log p_v}{e_v(q_v^{f_v} - 1)} + \epsilon.$$

Thus by letting  $\epsilon \rightarrow 0$  we can generate a sequence of numbers (in fact, algebraic integers) with the limit infimum of the height bounded by the desired constant.  $\square$

## References

- [1] M. BAKER, *A lower bound for average values of dynamical Green's functions*, Math. Res. Lett., **13** (2006), 245–257.
- [2] M. BAKER AND R. RUMELY, *Potential theory and dynamics on the Berkovich projective line*, Mathematical Surveys and Monographs **159**, American Mathematical Society, Providence, RI, (2010).
- [3] E. BOMBIERI AND U. ZANNIER, *A note on heights in certain infinite extensions of  $\mathbb{Q}$* , Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **12** (2001), 5–14.
- [4] V. FLAMMANG, *Two new points in the spectrum of the absolute Mahler measure of totally positive algebraic integers*, Math. Comp. **65** (1996), 307–311.
- [5] R. RUMELY, *The Fekete-Szegő theorem with splitting conditions. I*, Acta Arith. **93** (2000), 99–116.
- [6] R. RUMELY, *The Fekete-Szegő theorem with splitting conditions. II*, Acta Arith. **103** (2002), 347–410.
- [7] R. RUMELY, *Capacity theory on algebraic curves*, Lecture Notes in Mathematics **1378**, Springer-Verlag, Berlin, (1989).
- [8] C. J. SMYTH, *On the measure of totally real algebraic integers*, J. Austral. Math. Soc. Ser. A **30** (1980/81), 137–149.
- [9] C. J. SMYTH, *On the measure of totally real algebraic integers. II*, Math. Comp. **37** (1981), 205–208.

Paul FILI  
Department of Mathematics  
University of Rochester, Rochester, NY 14627  
E-mail: fili@post.harvard.edu